

# **Ethnobotanical Study on Assessment of Practice on Traditional Plant Medicine Use Among People of Wonchi District, Central Ethiopia**

Gadisa Melkamu \*

*Department of biology, Universty of Gonder, Ethiopia*

*Email: gadisamelkamu2010@gmail.com*

## **Abstract**

Ethiopia is a country characterized by a wide range of climate and ecological condition which helped to have high diversity of medicinal plants and up to 80% of the population use traditional medicine for primary health care. Studies on the current practices of the communities in the area are vital to document the basic information in these regars. Therefore, this study was aimed in assessing the practices of traditional plant medicine use among communities of Wonchi District.

**Keywords:** Indiginous practice; Local people; Medicinal plants; Traditional Medicine; Wonch district.

## **Results**

The most widely used route of administration in the study area was oral accounted for [56.67%] followed by dermal [29.63%] This is the reason that oral and dermal routes permit rapid physiological reaction of the prepared medicines with the pathogens and increase its curative power. The prepared traditional medicines were applied in a number of ways, among which drinking [37.57%], creaming [16.76%], and eating [10.40%] were mentioned frequently.

## **1. Introduction**

According to [83] consultation of medicinal practitioners is very helpful for the development and incorporation of useful approaches in planning and budgeting system for health care provision of most developing nations and indigenou communities.

---

\* Corresponding author.

In Africa, traditional medicine plays a central role in health care needs of rural people and urban poor. Here, it is said that, this situation would remain so long as modern medicine continues to be unable to meet the health care of the people of the continent effectively [41]. The value and role of this health care system will not diminish in the future, because they are both culturally viable and expected to remain affordable, while the modern health care service is both limited and expensive [83]. Indigenous traditional medicinal practices were carried out essentially based on private practice, i.e. private agreement between consenting parties, and the knowledge of traditional practice in most cases has descended through oral folk lore [6]. The secret information retained by traditional healers is relatively less susceptible to distortion but less accessible to the public. However, the knowledge is dynamic as the practitioners make every effort to widen their scope by reciprocal exchange of limited information with each other [16]. Incomplete coverage of modern medical system, shortage of pharmaceuticals and unaffordable prices of modern drugs, make the majority of Ethiopian still to depend on traditional plant medicines [17,71]. Hence the present study was initiated to investigate the indigenous practice of traditional plant medicine use among local communities of Wonch District, Western Ethiopia.


### ***1.1.Statement of the Problem***

Traditional medicine is an ancient form of health care practices long before appearance of scientific medicine which have played and continue to have important role in providing curative services to very large number of people particularly in the rural areas of almost all countries of Africa [35]. It is the culture of many people because of its accessibility to the people even in most remote areas particularly in the community where care is given at low cost to patients in their home. Most people have good attitude towards traditional plant medicine, although it is not always the best form of health care system [35]. In many parts of Ethiopia, considerable numbers of researches have been done on those practice of traditional plant medicine [5]. Like in other parts of country, in the current study area, the knowledge on medicinal plants depth and width become lesser and lesser due to its secrecy, unwillingness of young generation to gain the knowledge, influence of modern education, religious and awareness factors, which all results in gradual disappearance of indigenous knowledge on medicinal plants [Researcher long term direct observation]. But there was no much formal research work that had been done on the indigenous practice of traditional plant medicine in the study area. Therefore, this study was aimed to document the traditional medicinal plant species practices in the study area.

## **2. Materials and method**

### ***2.1. Descriptions of the Study area and Location***

Wonchi District is one of the Districts in the Southwest Shoa Zone, Oromia Region, Ethiopia, which is located 124 km away from southwest of Addis Ababa with the area coverage of 460,516 hectare and the altitude range between 1798m to 2118m above sea level. The administrative center of Wonchi is Chitu and it has beautyfull Creator Lake known as Wonchi Lake from which the district has got its name. As a result many tourists from inside and outside visit this natural lake every year and it is source of income for the country [wonch district health office report,2019].




**Figure 1:** Map of the Study Area.

## 2.2. Population

Demographically the district has a population of 119, 736 with almost equal gender ratio of 49.8% male and 50.2% female. The average family size is 6 and the average number of children perhousehold was nearly 4 indicating that it is found to focus of development intervention addressing child wellbeing to bring real development in the community. Religion wise, Orthodox constitute 58.9%, Protestants 39.6% and Muslims constitute 1.3% while the ethnic group composition, as per the Terminal evaluation findings of 2013, more than 99% are Oromo, the remaining being Amhara, Gurage and others [CSA, 2007].

## 2.3. Climate

Ecologically the district is divided in to dega or high land [40%] and woina dega or mini land [60%]. The mean annual rain fall of the area ranges from 1650-1800mm with annual temperature range of 10-30<sup>o</sup>c and mean average of 19.6 <sup>o</sup>c. The study area had 28.7<sup>o</sup>c annual mean maximum and, 19.6<sup>o</sup>c annual mean minimum temperature. The annual mean maximum and minimum temperature were recorded in March and November respectively. The highest rainfall distributions occur from June to September [Figure 2].



**Figure 2:** Climadiagram of the study area from 2006-2017.

#### 2.4. Land use types

Out of the total areas of the District, 82% is cultivated land, 11.7% grazing land, 8.9% covered by natural forest, 1.03% is water body while others is 18.6% [World Vision Ethiopia Wonchi area Office report, 2018].

#### 2.5. Vegetation of the study area

Due to variation in altitude and topographical features, the wonchi district vegetation shows three different zones, namely: Afromontane forest, sub alpine and afroalpine] vegetation [92,37]. The common plant species of the study area include: *Achyranthes aspera*, *Albizia schimperiana*, *Alchemilla pedata*, *Apodytes dimidiata*, *Brucea antidysenterica*, *Dombeya torrida*, *Embelia schimperi*, *Erica arborea*, *Festuca gilbertiana*, *Lobelia rhynchopetalum*, *Hagenia abyssinica*, *Hypericum revolutum*, *Jasminum abyssinicum*, *Juniperus procera*, *Kniphofia foliosa*, *Lobelia giberroa*, *Maytenus arbutifolia*, *Milletia ferruginea*, *Nuxia congesta*, *Olea capensis*, *Olea europaea subsp. caspidata*, *Papaneasimensis*, *Pittosporum viridiflorum*, *Prunus africana*, *Phytolacadicandra*, *Salix subserrata*, *Schefflera abyssinica*, *Thymus schimperi* and *Zehneria scabra* Vegetation.

#### 2.6. Study Design

Field survey design was employed together information on the indigenous knowledge, attitude and practice of traditional plant medicine of the local people in the study area. During the survey, both qualitative [none numerical] and quantitative [numerical] data were collected.

### **2.7. Reconnaissance Surveys**

Preliminary survey was conducted from march 20- 25, 2020. During the preliminary survey general information about the study area were gathered. Based on the information sampling technique, Sampled Kebeles, number of informants and study sites were determined.

### **2.8. Study Site Selection**

From a total 23 Kebeles in the District, nine study Kebeles were selected purposively based on availability of key informants following the recommendation of government officials, stakeholders, and religious leaders during reconnaissance survey. The sampled Kebeles are [Belbela,Dimtu,Fite ,Haro wanch,Kurfo gute,Lemen meta hora,Miti welga,Sonkole kake,Waldo telfa ].

### **2.9. Informant Selection**

A total of 198 informants were selected. From these 27 were key informants [3 informants per Kebele] which were selected purposively and 171 [19 per Kebele] of them were general informants which were selected randomly [simple random sampling technique following lottery method]. Age range of informants selected for the study were from 20 to 80 who lived 5 year and above in the study area. According to [42],the size of the sample depends on the available fund, time and other reasons and not necessarily depends on total population.

### **2.10. Data Collection Method**

Semi-structured interview, observation and guided field walks with informants were employed to obtain ethnobotanical data as used by [38]. Interview was based on a checklist of questions prepared beforehand in English and translated to local languages [Afaan Oromo]. Information regarding indigenous practice of local community towards traditional plant medicine of healers was recorded at the spot. Guided field observation was made on the medicinal plants to check the availability of the plant in the area, to know the habit and habitat of the plant. Focus group discussion was also made to get more information on medicinal plants practice

### **2.11. Data Analyses**

Descriptive statistics were used to analyze the data on medicinal plants use and associated indigenous knowledge of local community, their attitude on traditional plant medicine use and medicinal plant used by traditional plant medicine healers of the study area. The results were displayed and summarized in tables and figures by using percentage, frequency and texts. The most useful information gathered on medicinal plants which were analyzed through the descriptive statistics include application, methods of preparation, route of application, disease treated, and parts used and the habit of the plant.

## **3. Results and discussion**

### **3.1 Socio-demographic characteristics of respondent's**

A total of 198 informants including 27 key informants were selected. As pointed out by Martin [1995], the selection of key informants is commonly systematic. Most of the respondents [77.77%] were males [Table 1]. The majority of respondent’s age range was from 40-60[51.5%]. Most of the participants [86.86%] were married [Table 1]. Almost all religious leader respondents were followers of Orthodox Christian. From all respondents 33.83% were able to read and write. Number of farmers’ respondents predominated [33.33%] other respondents while NGO workers are lower in number [5.05%] [Table 1].

**Table 1:** Socio-demographic characteristics of respondents in the study area.

No	Variables	Response option	Frequency	Percentage [%]
1.	Sex	Male	154	77.77
		Female	44	22.23
		Total	198	100
2.	Age	20-40	14	7.07
		41-60	102	51.51
		61-80	82	41.41
3.	Marital status	Single	12	6.06
		Married	171	86.86
		Windowed	15	7.57
4.	Religion	Christian	142	71.71
		Muslim	27	13.63
		Waqefata	25	12.62
		Others	4	2.02
5.	Education	Uneducated	41	20.70
		Able to read and write	67	33.83
		12 complete	10	5.05
		10 complete	38	19.19
		Diploma	39	19.69
		Degree	3	1.51
6.	Occupational status.	Farmers	66	33.33
		Merchants	25	12.62
		Government employer	36	18.18
		NGO worker	10	5.05
		Others	61	30.80

**Table 2:** Distribution of informant groups by number.

No	Respondents	Male	Female	Total	Percentage [%]
1	Farmers	26	7	33	16.66
2	Merchants	16	9	25	12.62
3	Religious leaders	26	6	32	16.16
4	Health care workers	26	10	36	18.18
5	Traditional plant medicine users	46	8	54	27.27
6	Traditional plant medicine healers	14	4	18	9.10
7	Total	154	44	198	100

### **3.2. Mode of preparation and route of administration**

Concerning the preparation of traditional medicine, the local people employed various methods of preparation of traditional medicines for different types of ailments. The most principal method of TMP preparation reported was in the form of crushing [20%] and the least was cooking [1.6%] [Table 8]. This might be the effective extraction of the plant gives immediate response for health problems when crushed or pounded to increase its curative potential. The result is consistent with the findings of [36] in which crushing is highly reported method of remedy preparation. But it disagrees with the report of [42] which revealed that squeezing is the most used preparation method.

The most widely used route of administration was oral accounted for [56.67%] followed by dermal [29.63%] [Table, 8]. This is the reason that oral and dermal routes permit rapid physiological reaction of the prepared medicines with the pathogens and increase its curative power [33]. These results are consistent with findings of various ethnobotanical researches elsewhere in Ethiopia and other countries such as that of [55,26,41,33,7,28,44, 36,61].

**Table 3:** Mode of preparation and route of administration.


<b>Forms of preparation</b>	<b>Total</b>	<b>% of total</b>	<b>Administration</b>	<b>Remedy counts</b>	<b>Percentage[%]</b>
Crushing	25	20	Oral	153	56.67
Pounding & mixing	25	18.4	Dermal	80	29.63
Pounding&powdering	23	17.6	Nasal	19	7.04
Squeezing	21	16	Optical	6	2.22
Chewing	18	14.4	Auricular	5	1.85
Pounding & squeezing	6	4	Nasal and oral	4	1.48
Decoction	5	4	Neck	3	1.11
Cooking	2	1.6			
<b>Total</b>	<b>125</b>	<b>100</b>		<b>270</b>	100

### **3.3. Ways of applications and dosage of plant remedies**

The prepared traditional medicines were applied in a number of ways, among which drinking [37.57%], creaming [16.76%], and eating [10.40%] were mentioned frequently [Figure 6]. This finding is consistent with the finding of [26,7] in which drinking accounted the largest percentage of remedy.

The dosage of medicine to be administered is given by estimating age, the physical condition of the patient and the severity of the diseases. Amounts to be administered is also estimated by the use of measurements such as length of a finger [for bark, root and stem length], pinch [for powdered plant material] different measuring

materials [e.g. spoon, coffee cup, tea cup and glass cups] and number count [for sap/extract drops, leaves, seeds, fruits, bulbs, rhizomes and flowers]. But these measurements are not accurate enough to determine the precise amount. Some of the medicinal preparations are reported to have adverse effects on the patients. Informants reported that *Hagenia abyssinica*, *Phytolacca dodecandra* and some others are found to have adverse side effects like stomach pain, vomiting and diarrhea. The informants recommended additives for some of these adverse side effects, such as drinking of milk and barley soup immediately after intake of medicinal plants [55,56,43]. This study agreed with study made by [3] in Kilte Awlaleo District, Eastern zone of Tigray region of Ethiopia [39] in Amaro district, southern nations and nationalities of Ethiopia showed no agreement in accurate measurement or unit used among informants.



**Figure 3:** Application ways of remedies for human and livestock ailment treatment.

### 3.4. Conditions of preparation of remedies

The results showed that majority of the remedies were prepared using fresh material 50, [53.76%], while 15 species [16.13%] were used in the dried form and 28 [30.11%] either fresh or dried. Similar studies were also conducted by [53,70] which showed that using fresh materials for different health problems is more than dry materials or dry or fresh. This could be due to the fact that the fresh materials did not lose their volatile bioactive chemicals like oils, which could deteriorate on drying.

### 3.5. Disease types and related medicinal plants in the study area

In the area a total of 57 ailment types [both human and livestock ailments] were recorded along with the medicinal plants. From these disease types, wound is the most frequently mentioned ailment type and it is claimed to be treated by many number [25 species] of medicinal plants. This is followed by Malaria and stomach ache which are claimed to be treated by 14 and 13 species respectively. While Abortion, back pain, bilharzia, ear defect, goiter, infertility, retained placenta and syphilis are claimed to be treated by only a single medicinal plant species [Appendix 7]. Paired-wise comparison analysis on six most important TMPs claimed to



treat wound was performed. The result showed that *Acacia abyssinica* is the most useful and effective plant to treat wound followed by *Kalanchoe petitiiana* while *Olea europaea* ranked sixth. [Table 9] Preference ranking was also made on other six TMPs which were mentioned to treat malaria [Table 10]. The result showed that *Vernonia amygdalina* is the most preferred species that ranked first followed by *Juniperus procera*. *Eucalyptus globulus* is the least preferred species followed by *Lepidium sativum* [Table 10]. All of the species particularly the top ranked ones by preference and pair wise needs special urgent conservation action and sustainable uses. In this regard the results agree with the findings of [7].

**Table 4:** Pared-wise comparison on six more mentioned medicinsal plants against wound.

Species name	Respondents [1-7]							Sum	Rank
	1	2	3	4	5	6	7		
<i>Acacia abyssinica</i>	3	5	2	2	5	3	4	24	1 <sup>st</sup>
<i>Kalanchoe petitiiana</i>	4	0	3	2	3	4	5	21	2 <sup>nd</sup>
<i>Asparagus africanus</i>	5	3	4	1	1	2	3	19	3 <sup>rd</sup>
<i>Euphorbia abyssinica</i>	2	4	1	4	0	3	2	16	4 <sup>th</sup>
<i>Rumex nervosus</i>	1	2	2	1	4	2	1	13	5 <sup>th</sup>
<i>Olea europaea</i>	0	1	3	5	2	1	0	12	6 <sup>th</sup>
Total	15	15	15	15	15	15	15	105	

**Table 5:** Preference ranking on six most frequently reported plants claimed to treat malaria.

Species name	Respondents							Sum	Rank
	1	2	3	4	5	6	7		
<i>Vernonia amygdalina</i>	6	2	4	5	1	2	6	26	1 <sup>st</sup>
<i>Juniperus procera</i>	5	3	3	2	4	4	4	25	2 <sup>nd</sup>
<i>Allium sativum</i>	4	6	5	1	2	3	3	24	3 <sup>rd</sup>
<i>Zingiber officinale</i>	3	5	4	2	6	1	2	23	4 <sup>th</sup>
<i>Lepidium sativum</i>	3	5	2	4	2	1	3	20	5 <sup>th</sup>
<i>Eucalyptus globulus</i>	2	4	1	3	5	1	2	18	6 <sup>th</sup>

### 3.6. Major human diseases in the study area

In the study area, a total of 44 diseases of humans recorded were treated with a total of 50 plant species, where one species can treat a single disease or a number of diseases [Appendex 5]. Similarly, one ailment can be treated with a combination of plant species or single plant. For example, wound is treated with 25 species of

plants, malaria and stomach-ache with 14 species each; body swelling and evil eye treated with 10 species each, tonsillitis with 9 species. Fibril illness, scabies [itches] and skin rash treated with 7 species each [Appendix 5]. Most of the reported medicinal plants were used to treat human ailments [Appendix 5]. This showed that, the people of the study area are more knowledgeable and give great attention about human ailments as compared to livestock diseases. Similar results were recorded by Seyoum Getaneh [2009] in Debre Libanos District, North Shewa Zone of Oromia Region, Ethiopia. Medicinal plants recorded in this study also used as remedies in other part of the country. For instance, 28 species were mentioned[53], 9 species[20], 10 species [2], 61 species in[26], 30 plant species[33], and 59 plant species [65].

### ***3.7. Livestock diseases in the study area***

In comparison to human diseases, livestock diseases were treated with a few number of plant species in the study area. A total of 13 livestock ailments were identified that were treated by traditional medicinal plants in the area [Appendix 8] Common diseases affecting livestock health in the study area were bloating which was treated by 10 species, anthrax and leech by 6 species each, ectoparasite [lice] by 5 species, rabies by 3 species, erythroblasts, horse disease, retained placenta and coccidiosis are treated by 2 species each and the remaining diseases are treated by 1 species each [Appendix 8] In addition, proper documentation and understanding of farmer's knowledge, attitude, and practices about the occurrence, cause, treatments, prevention and control of various ailments is important in designing and implementing successful livestock production [75].

### ***3.8. Threats and conservation of medicinal plants in the study area***

#### ***3.8.1. Threats to medicinal plants***

The causes of threats to medicinal plants in the study area were both natural and anthropogenic factors. The most dominant factors affecting the medicinal plants in the study area was agricultural land expansion [34.34%] followed by charcoal production [16.16%]. While, the least serious factor was wild fire [4.04%] and then overflowing [4.54%] [Table 11]. Similar problems were also emphasized by [27,82]. Moreover, the problems identified so far during the course of this study are almost similar to what other literature sources studied in many parts of the country have already stated [35,55,80]. The medicinal plants of Wonch district in general and particular are facing the same problem. The loss of medicinal plants associated with the missing advantages gained from medicinal plants and indigenous knowledge associated with plants [69]. This is observed in wonch district as collection and search for some medicinal plants like *Cordia africana*, *Ekebergia capensis* and *Thalictrum rhyngocarpum* need longer time distance from their residence. Similar findings were also reported in Ethiopia [26,7,36] that showed need for agricultural land and for other uses severely threatened plant species in general and medicinal plants in particular.

Merchants, health care workers and other members of society obtained charcoal and timber from *Acacia abyssinica* and *Cordia africana* mature plants were recorded in the area indicating over exploitation. [7] argue that quite simply, mature seed producing tree that are the backbone of the population will die and are not replaced and ultimately the resource base on which culturally values are built will disappear because of over

harvesting.

Individual farmers in the area as observed during the study penetrated the forest with their axes daily. Here, the scenario is people need plants for their daily life activity i.e.as source of house hold tools, charcoal, furniture, agricultural implements. Thus, those multi-purpose species are on front line to be affected by these activities.

**Table 6:** Factors affecting Medicinal plants in the study area.

Variable	Factors	Frequency	Percentage [%]
Threats to conservation of medicinal plants	Agricultural land expansion	68	34.34
	Fire wood	23	11.61
	Charcoal	32	16.16
	Timber production	17	8.58
	Construction wood	21	10.60
	Medicinal plant trade	10	5.05
	Drought	10	5.05
	Over flooding	9	4.54
	Wild fire	8	4.04
Total		198	100

### 3.9. Conservation of medicinal plants and associated knowledge in the study area

Local people of the area know the importance of conserving the plants in both ex-situ and in-situ conservation methods. For instance, some people have started conserving the plants in fenced/protected pasture land [18.62%]; in different worship areas [churches, mosques] [21.49%], in their farms [18.62%], field/farm margins and around their home gardens [18.58%] and live fences of the famers [20.20%] [Table 12]. [59,36] have also reported that different worship areas are conservation sites for remnant vegetation in general and medicinal plants in particular. For instance, medicinal plants like *Juniperus procera*, *Olea europaea* subsp.cuspidata and *Euphorbia abyssinica* are found in church forest and also plants like *Hagenia abyssinica*, *Ocimum urticifolium* and *Ruta chalepensis* are found in the majority of home gardens in the study area, as they need these plants in their daily life as spices, medicine or for other values. Plants such as *Acacia abyssinica* and *Cordia africana* are also left as remnants of forest in the agricultural field due to their uses as timber source, for construction and fuel wood. Many medicinal plant species were also reported to be rare. Some of these local names are BOODAA WALEENSSUU [meaning plain land of *Erythrina brucei*], BARAA CALALQAA [meaning valley of *Apodytes dimidiata*], KARREE BAROODDOO [meaning hilly slope of *Myrica salicifolia*], and GULLUUGURRAA [meaning mountainous slop of *Prunus africana*]. What then ethno botanists have to learn from such evidences should be the point of focus. Such local clues could be good contributors for

designing ecosystem/habitat conservation, rehabilitation and resilience of species in their wild state where they are best adapted. These need an urgent attention to conserve such resources in order to optimize their use in the primary health care system. Some studies have shown that most of the medicinal plants used in Ethiopia are harvested from the wild [50, 88].

**Table 7:** Indigenous knowledge of local community towards medicinal plants conservation.

Variables		Frequency	Percentage [%]
Knowledge on the importance of medicinal plant conservation	Good	101	51.01
	Not good	97	48.99
Types of conservation	Total	198	100
	On worship areas	42	21.49
	On protected pasture	36	18.62
	In their farms	43	21.11
	In home gardens	37	18.58
	In live fences of the farmers	40	20.20
Total		198	100

#### 4. Conclusion and recommendations

##### 4.1. Conclusion

A study on medicinal plant indigenous knowledge, attitude and practice in the area revealed that the community use medicinal plants for maintaining their primary health care. From the study it can be said that the different segment of the community in the study area are in different level of knowledge with regard to traditional plant medicine use, i.e. difference in age, sex, work and education level has impact on the knowledge of the use of traditional plant medicine. In addition from the result of the study it can be concluded that there are considerable number of community members which do have negative attitude towards use of traditional plant medicines specially educated and youngsters are developing negative attitudes. Moreover, the result of the study revealed that, though negative attitude towards traditional plant medicine is believed to be increasing from time to time, still the community is extensively practicing the use of traditional plant medicines. The ethnobotanical study of medicinal plants in the study area showed that medicinal plants are used by a large member of the population and it is the most important means of treating some common human and livestock ailments. Most medicinal plants collected and identified were herbs and all plant parts were used for preparation of remedies. However, the use of medicinal plants for multiple purposes is leading to depletion in an alarming rate. This is

worthy because of some of the uses [Agricultural expansion, firewood, construction, forage, charcoal.] are the major destructive. Threats that erode indigenous knowledge usually comes from secrecy, oral-based knowledge transfer, the unwillingness of young generation to gain the knowledge, unavailability of the species, the influence of modern education and awareness factors are the major ones. The results of this study also showed that cultivation of plant species in and around home gardens for different purposes have great contribution to the conservation of medicinal plants and the associated knowledge.

#### **4.2. Recommendations**

Based on the results of the study, the following recommendations are forwarded.

- ❖ Integrated conservation and management program on medicinal plants focused on awareness development and active involvement of local community, governmental and non governmental bodies shall be practiced in the district.
- ❖ Young generation needs raising awareness to avoid negative impacts on the medicinal plants and associated knowledge in the area, hence, documentation of the medicinal plants of the area needs to be continued.
- ❖ Avoid uprooting of the plant species for medicinal purpose particularly before its flowering, fruiting and/seeding. If possible, it is better to use other parts of the medicinal plants such as leaves instead of root to protect them from the risk of extinction and endangering the species by collecting the roots or barks of the plants.
- ❖ Establishing traditional healers associations by providing supports like land, fund and assistances for cultivations of medicinal plants in the district would help to conserve medicinal plants.
- ❖ The societies have no good awareness with tradition plant medicine healers. So that all stakeholders should work together to change the situation and to benefit from traditional plant medicine.
- ❖ The government should create possible conditions and include to the teaching curricula about traditional plant medicine use
- ❖ To change the attitude of the society any concerned body should give trainings, seminars about traditional plant medicine use.
- ❖ The government and other officials should recognize the use of traditional plant medicine and also the healers of traditional plant medicine need any supports from concerned bodies.
- ❖ The insights of religious institution and health care institution should be positive and work together with traditional plant medicine.
- ❖ The user's negative attitude should be changed in to positive and the lack of knowledge about traditional plant medicine use also should be changed by giving training to them and through creating awareness. All stakeholders should develop positive attitude for traditional plant medicine healers. The healers of traditional plant medicine should use appropriate measurements to give the medicine for users.

#### **Abbreviations**

IK; indigenousknowledge, MM; modern medicine, MP; medicinalplants, TGPU; Traditional medicinal plant

use

## **Declarations**

### **Ethical approval**

Written ethical clearance was obtained from the research and ethical committee of the department of biology university of Gonder. A formal letter was written to Wonchi District Health and Agricultural Office and each Kebele administration to conduct the study. Written informed consent was sought and obtained from every participant who decided to take part in the study. They were assured about the confidentiality of their responses.

### **Consent for publication**

Not applicable

### **Availability of data and materials**

The datasets used and/or analysed during the current study are available from the author for reasonable request.

### **Completing interests**

The author declares that they have no financial and non-financial competing interests.

### **Author contributions**

GM was involved in the conception, design, analysis, interpretation, report and manuscript writing.

### **Acknowledgements**

I extend my deepest gratitude to those who participated in the study for their time to provide relevant information. I wish to extend my thanks to data collectors and supervisors. I am also indebted to all those who apply their effort in the process of this study. Finally, I am thankful to the University of Gonder for their financial support provided.

### **References**

- [1]. Abebe Demissie [2001]. Biodiversity conservation of medicinal plants, problems and prospects. In conservation and sustainable use of medicinal plants in Ethiopia, pp. 198-203.
- [2]. Abiyot Berhanu, Zemedu Asfaw and Ensermu Kelbessa [2006]. Ethnobotany of plants used as insecticides, repellents and anti-malarial agents in Jabitehnon District, West Gojjam. *Ethiop. J. Sci.*, **29**[1]: 87-92.
- [3]. Abraha Teklay, Balcha Abera and Mirutse Giday [2013]. An ethnobotanical study of medicinal plants used in Kilege District, Tigray Region of Ethiopia. *Journal of Ethnobiology and*

Ethnomedicine, **9**:65.

- [4]. Amare Getahun [1976]. Some common medicinal and poisonous plants in Ethiopia in folk medicine. Addis Ababa University, Ethiopia.
- [5]. Aschalew Lata and Takele Etana ,[2014].Assessment of knowledge, attitude and practice on traditional medicine in lag hare dire dawatown,Addisababa:Ethiopia.
- [6]. Asfaw Debela, Dawit Abebe and Kelbessa Urga [1999]. An overview of traditional medicine in Ethiopia: Prospective and Development Efforts. In: [Tami rat Ejigu, ed.] Ethiopian.
- [7]. Balick, m.j.and cox,P.A.[1996].plants,people and culture;Science of ethnobotany.Newyork,USA. Bahailu Etana [2010]. Ethnobotanical Study of Traditional Medicinal Plants of GomaWereda Jima Zone of Oromia Region, Ethiopia: M.Sc Thesis.Addis Ababa University,Addis Ababa.
- [8]. Bannerman, R. H. [1993]. Traditional Medicine and Health Care Coverage. World Health Organization, Geneva.
- [9]. Bayaferers,T[2000].A Floristic analysis and ethno botany study of seem wet land of chefa Area, south wello, Ethiopia.M.sc.Thesis, Addis Ababa University.
- [10]. Berhane Kidane, Vander Maesen, L.J.G., Van An del, T. and Zemedede Asfaw [2014].Ethnoveterinary medicinal plants used by the Maale and ari ethnic communities in Southern Ethiopia. Journal of Ethnopharmacology 153: 274-282.
- [11]. Berhe, C., Mengistu, B., W/Aregay, G., 1995. Women Lead in Protecting Food Germ plasm and Herbs for Health in Ethiopia [unpublished]. Report Submitted to Earth Care Africa, Nairobi, Kenya
- [12]. Buragohain,J.,[2011].Ethno medical plants used by the ethnic communities of Tinsukia district Assam, India. Recent Research in science and Technology 3[9], 31-42.
- [13]. Cotton, C.M. [1996]. Ethno botany: Principles and Applications. John Wiley and Sons Ltd.,Chichester, England, pp. 347-374.
- [14]. Cunningham, A.B. [1996]. People, Park and Plants use recommendations for multiple use zones and development alternatives around Bwindi: Impenetrable National Park, Uganda.In: People and Plants: Working Paper 4, pp.18-25. UNESCO, Paris.
- [15]. Daniel [1999].Biostatistics, A foundation for analysis in the health science 7th edition; New York John Wiley.
- [16]. DawitAbebe, [1986]. Traditional Medicine In Ethiopia. The attempt being made to promote it for Effective and letter utilization. SINET Ethiopian Journal of science pp., 61-69.Addis Ababa, Ethiopia.
- [17]. Dawit and AheduAyelu. [1993]. Medicinal plants and Enigmatic Health practice of Northern Ethiopian. Publishing house Addis Ababa, Ethiopia.
- [18]. Dawit Abebeb, Asfaw Debela and KelbessaUrga [2005], Medicinal plants and other useful.
- [19]. Dawit Abebe [1986]. Traditional medicine in Ethiopia: The Attempts being made to promote it for effective and better Utilization. SINET: Ethiop. J.Sci. [Supp.]: 62-69.
- [20]. Debela Hunde, Zemedede Asfaw and Ensermu Kelbessa [2004]. Use and Management of Ethnoveterinary Medicinal Plants by Indigenous People in Boosat, Welenchiti area. Ethiopia. J. Biol. Sci., 3 [2]: 113-132.
- [21]. Debela Hunde, MesfinTadesse and YihenewGirma.[2005] Survey of Medicinal plants used to Treated human disease in sekachekorsa, Jimma Zone, Ethiopia,J,Heath Sci.15:89-106.

- [22]. Debela H[2001].Use and management of traditional medicinal plant by indiginous people ofboosat distirct,wolencht area:An ethno botanical approach.M.SC.thesis,Addis Ababa university p.48.
- [23]. Deilnesaw yehualawu, Haile Yenegere and TizazuGebrie [2005], ethno botanical study of Traditional medicinal plants ,Haramaya Ethiopia.
- [24]. Demel Teketay [2001]. Deforestation, Wood Famine and Environmental Degradation in Ethiopia's Highland Ecosystems: Urgent Need for Action. *Northeast African Studies*, [New Series]. 8 [1]: 53-76
- [25]. Endashaw Bekele [2007]. Study on Actual Situation of Medicinal Plants in Ethiopia. Moringa stenopetala is a perennial tree.
- [26]. Endalew Amenu [2007]. Use and Management of Medicinal Plants by indigenous People of Ejaji Area [ChelyaWereda] West Shewa, Ethiopia: An Ethnobotanical Approach, M.ScThesis.Addis Ababa University, Addis Ababa.
- [27]. Ensemu Kelbessa, Sebsebe Demissew, Zerihun Woldu and Edwards, S. [1992]. Some threatened Endemic Plants of Ethiopia. In: [S. Edwards and Zemedede Asfaw eds.]. *The Status of Some Plants in Parts of Tropical Africa. NAPRECA, Series No.2, BOTANY 200: East and Central Africa*.Pp. 35-55.
- [28]. Ermias Lulekal, Zemedede Asfaw, Ensemu Kelbessa and Damme, P.V. [2013]. Ethnomedicinal Study of plants used for human ailments in Ankober District, North Shewa Zone, Amhara Region, Ethiopia. *Journal of Ethnobiology and Ethnomedicine* 9:63.
- [29]. Eskedar Abebe [2011].Ethno botanical study on medicinal plants used by local communities in Debarke distirct, North Gondar zone, Amhara regional state.
- [30]. Etana T[2007].Use and conservation of traditional medicinal plants by indiginous people in Gimbidistirct, westernwollega, Ethiopia,M.sc thesis,AAU,Ethiopia.
- [31]. Fassil, H., 2003. Ethiopia: a qualitative understanding of local traditional knowledge and medicinal plant use. *IK Notes* 52, 1–4
- [32]. Gedif, T., Hahn, H., 2003. The use of medicinal plants in self-care in rural central Ethiopia. *Journal of Ethnopharmacology* 87, 155–161
- [33]. Fisseha Mesfin, Sebsebe Demissew and Tilahun Teklehymanot [2009]. An ethnobotanical study of medicinal plants in Wonago district, SNNPR, Ethiopia. *Journal of Ethnobiology and Ethno medicine* 5: 28. <http://www.Ethnobiomed.com/content/5/1/28>. Accessed on December 26, 2018.
- [34]. Fullas F [2001] .Ethiopian Traditional Medicine: Common Medicinal Plants in Perspective Edition. FekaduFullas: Iowa.
- [35]. Getachew Addis, Dawit Abebe and Kelbessa Urga [2001]. A Survey of Traditional Medicinal Plants in Shirka Ditirct, Arsi Zone, Ethiopia. *Pharmaceutical Journal* 19:30-47.
- [36]. Getnet Chekole [2015].Ethnobotanical Study of Medicinal Plants in the environs of Tara-Gedam and Amba Remnant Forests in Libo Kemkem District. *Journal of Ethnobiology and Ethnomedicine*.
- [37]. Getinet Masresha [2014]. Diversity, Structure and Regeneration Status of Vegetation in Simien Mountains National Park, Northern Ethiopia: PhD. Dissertation. Addis Ababa University, Addis Ababa.
- [38]. Getu Alemayehu [2010]. Ethnobotanical Study of Medicinal Plants used by Local communities Of Minjar-Shenkora District, North Shewa Zone of Amhara Region, Ethiopia. *Journal of Medicinal Plants Studies*, 3[6].



- [39]. Getu Alemayehu [2017]. Ethnobotanical study of plant diversity and ethnobotany of medicinal and wild edible plants in amaro district of southern nations, nationalities and peoples region and gelana district of oromia region, southern Ethiopia.
- [40]. Haile Yineger [2005]. A Study on the Ethno botany of Medicinal Plants and Floristic Composition of the dry afro Montana Forest at Bale Mountains National Park, Ethiopia: M.Sc. Thesis. Addis Ababa University West. *International Journal of Medicine and medicinal Sciences Health spa &Resort. Thailand.*
- [41]. Jansen, P.C.M. [1981]. Spices, Condiments and Medicinal plants in Ethiopia, their Taxonomy and Agricultural Significance. Center for Agricultural Publishing and Documentation, Wageningen, Netherlands. Pp 327.
- [42]. Jarsso Belay [2016]. Ethnobotanical Study of Traditional Medicinal Plants used by Indigenous People of Jiggiga district, Somali Regional State, Ethiopia: MSc.Thesis.Haramaya University, Haramaya.
- [43]. Kebu Balemie, Ensermu Kelbessa and Zemedede Asfaw [2004]. Indigenous Medicinal Plant Utilization, Management and Threats in Fentalle Area, Eastern Shewa, Ethiopia. *Ethiopian Journal of Science* 3 [1]: 37-58.
- [44]. Ketema Tolossa, Etana Debela, Spiridoula Athanasiadou, Adugna Tolera, Gebeyehu Ganga and Jos GM Houdijk [2013]. Ethno-medicinal study of plants used for treatment of human and livestock ailments by traditional healers in South Omo, Southern Ethiopia. *Journal of Ethnobiology and Ethnomedicine* 9:32.
- [45]. Konno B. [2004]. International of traditional medicine with modern medicine. *EHNRS. Addis Abebe. app, 3-9.*
- [46]. Lai, PK., and Roy, J. [2004]. Antimicrobial and chemo preventive properties of herbs and Spices. *Curr. Med. Chem.* 11 [11]: 1451–60.
- [47]. Lulekal, E., Kelbessa, E., Bekele, T., Yineger, H., 2008. An ethnobotanical study of medicinal plants in Mana Angetu District, southwestern Ethiopia. *Journal of Ethnobiology and Ethnomedicine* 4, 10.
- [48]. Manandhar, N.P. [1995]. A survey of medicinal plants of Jajarkot District, Nepal. *Journal of Ethnopharmacology* 48[1]:1-6.
- [49]. Martin, G. J. [1995]. *Ethnobotany: A method Manual. A 'People and Plants' Conservation Manual.* Chapman and Hall, London, pp. 268.
- [50]. Mesfin Tadesse and Sebsebe Demissew [1992]. Medicinal Ethiopian plants inventory, identification and classification. In: *Plants used in African Traditional medicine as Practiced in Ethiopia and Uganda, East Africa.* [Edwards, S. and Zemedede Asfaw, ed]. Monograph Series. No. 5:1-19. Addis Ababa.
- [51]. Mirutse G. [1999]. An Ethnobotanical study of medicinal plants used by the people in Ethiopia.
- [52]. Martin, G.J. [1995]. *Ethno botany: A method Manual.* Chapman and Hall, London. Pp. 265-270. ce. 30:18-22.
- [53]. Mengistu Gebrehiwot [2010]. An Ethnobotanical Study of Medicinal Plants in Seru district, Arsi zone of Oromia Region, Ethiopia: M.Sc. Thesis. Addis Ababa University, Addis Ababa.
- [54]. Mesfin Tadesse [1986]. Some Medicinal Plants of Central Shewa and South Western Ethiopia. *SINET: Ethiop. J. Sci.*, 9: 143-167..

- [55]. Mirutse Giday and Gobena Ameni [2003]. An Ethnobotanical Survey on Plants of Veterinary Importance in two Woredas of Southern Tigray, Northern Ethiopia. *SINET: Ethiopian J. of Sci.*, **26**:123-136.
- [56]. Mirutse Giday, Zemedede Asfaw, Thomas Equist and Zerihun Woldu [2003]. An Ethnobotanical Study of Medicinal Plants used by the Zay people in Ethiopia. *Journal of Ethnopharmacology* 85:43-52.
- [57]. Mulugeta Kuma [2014]. Use and Management of Medicinal Plants by Indigenous People of Jima Rare District in Oromia Region, Ethiopia: M.Sc. Thesis. Haramaya University, Haramaya.
- [58]. Mutabazi [2008]. Traditional knowledge and imperialistic notion. *Rabist.com*, 2015.
- [59]. Nigussie Amsalu [2010]. An Ethnobotanical Study of Medicinal Plants in Farta Woreda, South Gondar Zone of Amhara Region, Ethiopia: M.Sc Thesis. Addis Ababa University, Addis Ababa.
- [60]. Pharmacotherapy Group [2009]. Current Trends in Ethnobotany. *Tropical Journal of Pharmaceutical Research* 8[4]: 295-296.
- [61]. Patience, T., Esezah, K., Mukadasi B., Justine N., Maud K., Patrick M. and James K. [2016]. Ethnobotanical survey of medicinal plant species used by communities around Mabira Central Forest Reserve, Uganda. *Journal of Ethnobiology and Ethnomedicine* 12:5.
- [62]. Quanash, N. [1998]. Bicultural diversity and integrated health care in Madagascar. *Nature and Resource*.
- [63]. Robert and Henry, [2000] *Medicinal Indian* 2<sup>nd</sup> edition Asiatic publishing house. New Delhi India.
- Sherman, P. and Hash, G.A. [2001]. Why vegetable recipes are not very spicy. *Evol. and Hu Behav.* 22 [3]: 147–163.
- [64]. Sebsebe Demissew and Ermias Dagne [2001]. Basic and applied research on medicinal plants of Ethiopia. In: Conservation and Sustainable Use of Medicinal Plants in Ethiopia, Proceeding of The National Work Shop on Biodiversity and Sustainable use of Medicinal Plants In Ethiopia, 28 April-01 May 1998, pp 29-33. [Medhin Zewdu and Abebe Demissei eds.], IBCR, AA.
- [65]. Seyoum Getaneh [2009]. Ethnobotanical study of Medicinal Plants in Debre-Libanos district, North Shewa Zone of Oromia Region, Ethiopia M.Sc. Thesis, Addis Abeba University, pp. 94.
- [66]. Sintayehu Tamene [2011]. An Ethnobotanical Study Of Medicinal Plants In Wondo Genet Natural Forest and Adjacent Kebeles, Sidama Zone, SNNP Region, Ethiopia. MSc. Thesis. Addis Ababa University, Addis Ababa, Ethiopia.
- [67]. Sumner, J. [2000]. *The Natural History of Medicinal Plants*. Timber Press. p. 16.
- [68]. Stepp, J. [2004]. The role of weeds as sources of pharmaceuticals. *Journal of Ethno pharmacology*, 92 [2–3]: 163–166.
- [69]. Sofowara, A. [1982]. *Medicinal plants and traditional medicine in Africa*. John William and sons. New York, pp. 225-256.
- [70]. Tadesse Beyene [2015]. Ethnobotany of Medicinal Plants in Erob and Gulomahda Districts, eastern Zone of Tigray Region, Ethiopia: PhD. Dissertation. Addis Ababa University, Addis Ababa.
- [71]. Teferi Gedife and Hahn, H. [2003]. The use of Medicinal Plants in self-care in rural central Ethiopia. *Journal of Ethnopharmacology* 87: 155-161.
- [72]. Tena Regassa [2016]. Vascular Plant Diversity and Ethnobotanical Study of Medicinal and wild edible Plants in Jibat, Gedo and Chilimo Forests, West Shewa Zone of Oromia region, Ethiopia: PhD. Dissertation. Addis Ababa University, Addis Ababa.

- [73]. Tewodros Tesfaye [2016]. Use and Management of Medicinal Plants by People of Melka Belo District, East Hararghe, Oromia Region, Ethiopia: M.Sc. Thesis.Haramaya University, Haramaya.
- [74]. Thomas, H. [1995]. Indigenous Knowledge, Emancipation and Alination. Journal of Knowledge transfer and utilization. 8[1]: 63-73. University of Washington.
- [75]. Tafesse Masfin and Mokonnen Lemma, [2001], plants in Ethiopia veterinary medicine. Their interaction with conventional drug.
- [76]. Teferi Flatie, Teferi Gedif, Kaleab Asres and Tsige Gebre-Mariam [2009] Ethnomedical survey of Berta ethnic group Assosa Zone, Benishangul-Gumuz regional state, mid-west Ethiopia. J. Ethnobiology and Ethnomedicine. 5:14
- [77]. Tesfaye A, Zemedede A [1999]. An Ethno-botanical Study of the Bertha People of the Benshangul Gumuz Region in Western Ethiopia. Program and Abstracts of the National Workshop. "Have We Valued Our Biodiversity?"
- [78]. Tesfaye Awas and Sebsebe Demissew [2009]. Ethnobotanical study of medicinal plants in Kafficho people, southwestern Ethiopia. In: Plant Diversity in Western Ethiopia: Ecology, Ethnobotany and Conservation. The University Foundation for Student Life [Sio]. University of Oslo, 2007. Norway. Pp 1-15."
- [79]. Tesfaye Hailemariam, Sebsebe Demissew and Zemedede Asfaw [2009]. An ethnobotanical study of medicinal plants used by local people in the lowlands of Konta special Wereda, Southern Nations, Nationalities and People Regional State, Ethiopia. Journal of Ethnobiology and Ethnomedicine, 6:25.
- [80]. Tilahun Teklehaymanot and Mirutse Giday [2007]. Ethnobotanical Study of Medicinal Plants used in the Zegie Peninsula, Northwestern Ethiopia. Journal of Ethnobiology and Ethno medicine. 3:12.
- [81]. Tizazu Gebre [2005]. An ethno botanical study of medicinal plants in konso special district, Southern nations, and nationalities and people's regional state, Ethiopia M.Sc. Thesis. Addis Ababa University, Addis Ababa.
- [82]. Vivero, J.L., Ensermu Kelbessa and Sebsebe Demissew [2005]. The Red List of Endemic Tree & Shrubs of Ethiopia and Eritrea. Published by Fauna and Flora International [FFI], Cambridge, UK.
- [83]. WHO [1979]. The promotion and development of traditional Medicine. World Health Organization. Technical Report Series 622, WHO, Geneva.
- [84]. WHO [1998]. Regulatory situation of herbal medicines: A World wide Review. Pp. 1-9. WHO/TRM/98.1, Geneva.
- [85]. WHO [2002]. Traditional medicine, growing needs and potential. Geneva. Zemedede Asfaw [2001], Conservation of traditional medicinal plants.
- [86]. WHO [2008]. Traditional medicine factsheet no. 134. December, 2008. <http://www.who.int/mediacentre/factsheets/fs123/en/> [Accessed,
- [87]. William, P.C. [2003]. Environmental Science: Global Concern. 11<sup>th</sup> ed. America, New York: McGraw-Hill.
- [88]. Zemedede Asfaw [2001]. The Role of Home garden in Production and Conservation of Medicinal Plants. In: Conservation and Sustainable Use of Medicinal plants in Ethiopia p. 76- 91, [Medhin Zewdu and Abebe Demissie Ed.]. Proceeding of the national workshop On Biodiversity Conservation and Sustainable use of medicinal plants in Ethiopia, 28 April – 01 May 1998, IBCR, Addis Ababa.

- [89]. Zelalem mengesha [2006]. Traditional vs. modern medicine from Jimma Town south east Ethiopia: publishing house Addis Ababa Ethiopia.
- [90]. Zewdie Kassa [2009]. An ethnobotanical study of medicinal plant biodiversity of trees and Shrubs in Jeldu Wereda, Western Shoa, Ethiopia. M.Sc. Thesis Addis Ababa University, Addis Ababa.
- [91]. Zerihun Woldu [1999]. Forest in the vegetation types of Ethiopia and their status in the geographical context. In: Edwards, S., Abebe Demissie, Taye Bekele and Haase, G. [eds.]. forest Genetic Resource Conservation: Principles, Strategies and Actions. Workshop proceedings. Institute of Biodiversity Conservation and Research, and GTZ, AA, Pp 1-41.

**Appendix 1:** Lists of plants collected from the study area [D=dry; F=fresh; DF= dry or fresh; Cl=climber; H=herb; Sh=shrub; T=tree]

**Table 13**

No	Scientific name	Family	Local name[oromoo]	Condition of plant	Plant habitat	Habit
1.	Acacia abyssinica Hochst. ex Benth	Fabaceae	Girar	F	wild	T
2.	Achyranthesaspera L.	Amaranthaceae	Talenj	F	wild	H
3.	Allium cepa L.	Alliaceae	Shinkurt Dimma	F	cultivated	H
4.	Allium sativum L.	Alliaceae	Shinkurt Addii	F	cultivated	H
5.	Aloe pubescens Reynolds	Aloaceae	Eret	F	wild	H
6.	Artemisia abyssinica Sch. Bip.ex A. Ri	Asteraceae	Harit	F	wild	H
7.	Asparagus africanus Lam.	Asparagaceae	Yesetkest	F	wild	Cl
8.	Brassica carinata A.Br.	Brassicaceae	Gomenzer	D	cultivated	H
9.	Brassica nigra L.	Brassicaceae	Senafich	D	cultivated	H
10.	Brucea antidysenterica Fresen.	Simaroubaceae	Waginos	DF	wild	Sh
11.	Buddleja polystachya Fresen.	Loganiaceae	Anfar	F	Both	Sh
12.	Calpurnia aurea [Ait.] Benth.	Fabaceae	Digita	DF	wild	Sh
13.	Capsicum annum L.	Solanaceae	Berbere	D	cultivated	H
14.	Carica papaya L.	Caricaceae	Papaya	F	cultivated	T
15.	Carissa spinarum L.	Apocynaceae	Agam	DF	wild	Sh
16.	Citrus limon [L.] Burm.f.	Rutaceae	Lomi	F	cultivated	Sh
17.	Clematis simensis Fresen.	Ranunculaceae	Azo hareg	DF	wild	Cl
18.	Clerodendrum myricoides [Hochst.] Vatke	Lamiaceae	Misirich	DF	wild	Sh
19.	Clutia abyssinica Jaub. & Spach	Euphorbiaceae	Fiyalefaj	F	wild	H
20.	Coffea Arabica L.	Rubiaceae	Buna	D	cultivated	Sh
21.	Cordia africana Lam.	Boraginaceae	Wanza	D	wild	T
22.	Coriandrum sativum L.	Apiaceae	Denbelal	D	cultivated	H
23.	Croton macrostchys Del.	Euphorbiaceae	Bisana	DF	wild	T

24.	<i>Cucumis ficifolius</i> A.Rich.	Cucurbitaceae	Ye medir enbuay	F	wild	Cl
25.	<i>Cucurbita pepo</i> L.	Cucurbitaceae	Duba	D	cultivated	Cl
26.	<i>Cupresses lusitanica</i> Mill.	Cupressaceae	Yeferenj tsid	F	Both	T
27.	<i>Cynoglossum coeruleum</i> [Hochst.ex.A.rich.]	Boraginaceae	Chigogit	F	wild	H
28.	<i>Cyphostemma adenocaula</i> [Steud. exA. Rich.]Des. ex Wild &Drum.	Vitaceae	Aserkush	F	wild	Cl
29.	<i>Datura stramonium</i> L.	Solanaceae	Astanagir	F	wild	H
30.	<i>Discopodium penninervum</i> Hochst.	Solanaceae	Almit	F	wild	Sh
31.	<i>Dodonaea angustifolia</i> L.	Sapindaceae	Kitkita	DF	wild	Sh
32.	<i>Dombeya torrida</i> [J. F. Gmel.] P. Bamps	Sterculiaceae	Wulkifa	DF	wild	T
33.	<i>Dovyalis abyssinica</i> [A.Rich.]Warb.	Flacourtiaceae	Koshim	F	Both	Sh
34.	<i>Echinops kebericho</i> Mesfin	Asteraceae	Kerebicho	D	wild	H
35.	<i>Embelia schimperi</i> Vatke	Myrsinaceae	Enkoko	DF	wild	Sh
36.	<i>Erica arborea</i> L.	Ericaceae	Asta	DF	wild	T
37.	<i>Eucalyptus globulus</i> Labill.	Myrtaceae	Bahirzaf Addi	DF	cultivated	T
38.	<i>Euphorbia abyssinica</i> . Gmel	Euphorbiaceae	Kulkual	DF	wild	T
39.	<i>Guizotia abyssinica</i> L.	Asteraceae	Nug	D	cultivated	H
40.	<i>Hagenia abyssinica</i> [Bruce] J. F. Gmel.	Rosaceae	Koso	DF	wild	T
41.	<i>Hypericum revolutum</i> Vahl	Hypericaceae	Amja	DF	wild	Sh
42.	<i>Jasminum abyssinicum</i> L.	Oleaceae	Tembebel	F	wild	Cl
43.	<i>Juniperus procera</i> Endl.	Cupressaceae	Ye abesha tsid	DF	wild	T
44.	<i>Justicia schimperiana</i> [Hochst. exNees] T. Anders	Acanthaceae	Simiza sensel	F	wild	Sh
45.	<i>Kalanchoe petitiiana</i> A. Rich.	Euphorbiaceae	Endawula	F	wild	H
46.	<i>Kosteletzkya begonifolia</i> Ulbr.	Malvaceae	Yemegerem	F	wild	H
47.	<i>Linum usitatissimum</i> L.	Lineaceae	Telba	D	cultivated	H
48.	<i>Lobelia rhyngopetalum</i> Hemsl.	Lobeliaceae	Jibira	D	wild	Sh
49.	<i>Lepidium sativum</i> L.	Brassicaceae	Feto	D	cultivated	H
50.	<i>Maesa lanceolata</i> Forssk.	Myrsinaceae	Shwaria[kelawa]	DF	wild	Sh
51.	<i>Melia azedarach</i> Forssk.	Meliaceae	Nim	DF	cultivated	T
52.	<i>Myrtus communis</i> L.	Myrtaceae	Ades	D	wild	Sh
53.	<i>Nicotiana tabacum</i> L.	Solanaceae	Timbaho	DF	cultivated	H
54.	<i>Nigella sativa</i> L.	Ranunculaceae	Tikur azmud	D	cultivated	H
55.	<i>Ocimum basilicum</i> L.	Lamiaceae	Zikakibe	F	cultivate	H
56.	<i>Ocimum lamiifolium</i> Hochst. ex Benth.	Lamiaceae	Damakese	F	Both	sh

57.	<i>Olea europaea</i> L. subsp. <i>cuspidata</i> [Wall. ex G. Don] Cif.	Oleaceae	Weyra	DF	wild	T
58.	<i>Opuntia ficus-indica</i> [L.] Miller	Cactaceae	Balas[Ashewa kulkual]	F	wild	Sh
59.	<i>Otostegia integrifolia</i> Benth.	Lamiaceae	Tunjit	D	wild	Sh
60.	<i>Osyris quadripartita</i> Decn.	Santalaceae	Keret	F	wild	Sh
61.	<i>Phytolacca dodecandra</i> L' Herit.	Phytolaccaceae	Endod	F	wild	Sh
62.	<i>Pittosporum viridiflorum</i> Sims	Pittosporaceae	Tebera	F	wild	T
63.	<i>Plantago lanceolata</i> L.	Plantaginaceae	Gortab [Wenberet]	F	wild	H
64.	<i>Prunus persica</i> [L.] Batsch	Rosaceae	Kok	F	cultivated	T
65.	<i>Pterolobium stellatum</i> [Fors]Brenan	Fabaceae	Kentafa	DF	wild	Sh
66.	<i>Rhamnus prinoides</i> L. Herit.	Rhamnaceae	Gesho	F	cultivated	Sh
67.	<i>Ricinus communis</i> L.	Euphorbiaceae	Gulo	DF	cultivated	Sh
68.	<i>Rosa abyssinica</i> Lindley	Rosaceae	Kega	F	wild	Sh
69.	<i>Rumex abyssinicus</i> Jacq.	Polygonaceae	Makmako	DF	wild	H
70.	<i>Rumex nepalensis</i> Spreng.	Polygonaceae	Lut	F	wild	H
71.	<i>Rumex nervosus</i> Vahl	Polygonaceae	Embacho	DF	wild	Sh
72.	<i>Ruta chalepensis</i> L.	Rutaceae	Tenadam	DF	cultivated	H
73.	<i>Salix macronata</i> Thunb.	Salicaceae	Lahaya	F	wild	T
74.	<i>Schinus molle</i> L.	Anacardiaceae	Kundoberbere	DF	cultivated	T
75.	<i>Sesamum indicum</i> L.	Pedaliaceae	Selit	D	cultivated	H
76.	<i>Sida schimperiana</i> Hochst. Ex A. Rich.	Malvaceae	Cifrig	DF	wild	H
77.	<i>Solanum adoense</i> [Hochst] ex A. Rich.	Solanaceae	Zerch Enbuay	DF	wild	Sh
78.	<i>Solanum incanum</i> L.	Solanaceae	Enbuay	DF	wild	Sh
79.	<i>Solanecio gigas</i> [Vatke.]C. Jeffery	Asteraceae	Yeshikoko gomen [Mogne qitel]	F	wild	Sh
80.	<i>Stephania abyssinica</i> [Dillo&A.Rich.]Walp.	Menispermaceae	Ye ayit hareg [Chewchawit]	F	wild	Cl
81.	<i>Syzgium aromaticum</i> L.	Myrtaceae	Kirunfu	D	cultivated	T
82.	<i>Tanacetum cinerariifolium</i> [Trev]. Sch. Bip.	Asteraceae	Kaba	F	wild	H
83.	<i>Trigonella foenum-Graecum</i> L.	Fabaceae	Abish	D	cultivated	H
84.	<i>Thymus schimperi</i> Ronniger	Lamiaceae	Tosign	DF	wild	H
85.	<i>Urera hypselodendron</i> [A. Rich.] Wedd.	Urticaceae	Lankuso	F	wild	Cl
86.	<i>Urtica simensis</i> Steudel	Urticaceae	Sama	F	wild	H
87.	<i>Verbena officinalis</i> L.	Verbenaceae	Atuch	F	wild	H
88.	<i>Verbascum sinaiticum</i> Benth.	Scrophulariaceae	Kutina[ye Ahya joro]	DF	wild	H
89.	<i>Vernonia amygdalina</i> Del.	Asteraceae	Gerawa	F	Both	T

90.	Vernonia hymenolepis A.Rich.	Asteraceae	Weynagift	F		Sh
91.	Zehneria scabra [Linn.f.] Sond.	Cucurbitaceae	Haregresal[Etse sabeq[NechHareg]	F	wild	Cl
92.	Zingiber officinale Roscoe	Zingiberaceae	Jinjible	F	cultivated	H
93.	Ziziphus spina- christi [L.] Desf.	Rhamnaceae	Gaba	F	Both	Sh

**Appendix 2:** List of medicinal plants used for both human [Hu] and livestock [Li] ailment treatment: scientific name; family; local name [A=Amharic; O=Afan Oromo] habit; parts used; disease treated; methods of preparation with dosage used and route of application.

**Key:**Habit [Ha.]: Herb [H] ;Shrub [Sh] ; Tree[T]; Climber [Cl.]; Epiphyte [Ep.]; Parts used [Bark, B; Latex, La; Root, R; Leaf, L; Fruit, Fu.; Flower, Fw.; Seed, Se.;Stem,St.;Sap,Sa; Bulb, Bu.; Root and leaf, LR.; Above ground , Ag; Leaf and seed, LSe; shoot, Sht.

**Table 14**

Scientific name	Family	Local Name	Ha	Pu	Use d for	Disease treated	Mode of preparation	Route
Acacia abyssinica Hochst. ex Benth.	Fabaceae	Girar [A] Laaftoo[O]	T	Sht	Hu	Tonsillitis	Fresh shoot is chewed and the juice is swallowed.	Oral
				Sht	Hu	Wound	Fresh shoot is pounded, squeezed and the sap is creamed on affected part	Dermal
Achyranthes aspera L.	Amarantaceae	Talenj[A] Darguu[O]	H	L	Both	Pneumonia	Fresh leaf are pounded, mixed with water and drunk in the Morning.	Oral
				L	Hu	Body swelling	Leaf is squeezed and creamed on swollen body.	Dermal
				L	Hu	Wound	Leaves are pounded and tied on the wound.	Dermal
				L	Hu	Nasal bleeding	Fresh leaf of Achyranthes aspera is squeezed and its juice is dropped in to the nostrils to stop bleeding.	Dermal
				R	Hu	Stomach-ache	The root of Achyranthes aspera is chewed and swallowed during feeling of ache.	Oral
Allium cepa L.	Alliaceae	Qey shenkurt[A] Qullubbii Diimaa[O]	H	Bu	Li	Leech	Fresh bulb is pounded, mixed with little water, decanted and poured through the nose.	Oral
					Hu	Asthma	Fresh bulb of Allium cepa is crushed with Zingiber officinale rhizome and boiled. Then drink the solution in the morning until recovery.	Oral
Allium sativum L.	Alliaceae	Nech shinkurt[A] Qullubbii adii[O]	H	Bu	Hu	Malaria	Fresh bulb of Allium sativum and rhizome of Zinger officinale are pounded together, mixed with honey and eaten.	Oral
				Bu	Hu	Evil eye	Bulb of Allium sativum is crushed together with rhizome of Zingiber officinale and Lepidium sativum, pasted with honey and 2 tea spoons is taken.	Oral
				Bu	Hu	Colds	The Bulb is pounded, mixed with honey and 2-3 teaspoon is eaten every day for three days.	Oral
Aloe pubescens Reynolds	Aloaceae	Eret [A] Argiisa[O]	H	La	Hu	Fire burn	The latex is painted on the wound	Dermal
				La	Hu	Ringworm	The latex of the young leaves is creamed on the affected area and repeated	Dermal



							every day until recovery.	
				La	Li	anthrax	Root of Aloe pubescens is crushed and mixed with cold water. Then two cup of tella is given to cattle.	Oral
Artemisia abyssinica Sch. Bip.ex A. Rich.	Asteraceae	Harit[A] Harrittaa[O]	H	L	Hu	Evil eye	Leaf concoction together with root of Echinops kebericho is added to a burning fire and smoked to the patient.	Nasal
				L	Hu	Malaria	Fresh leaf is crushed and pounded with water, filtered and drunk until recovery.	Oral
				L	Hu	Child Stomach-ache	Leaf is squeezed and ¼ of coffee cup juice is given to children	Oral
Asparagus africanus Lam.	Asparagaceae	Yesetkest[A] Sariitii[O]	Cl	R	Hu	Retained placenta	Fresh root is pounded; mixed with water and boiled, decanted, mixed with honey and drunk.	Oral
				R	Hu	Wound	Fresh root is pounded together with leaves of Dodonea angustifolia, mixed with butter and creamed on the wound.	Dermal
				L	Hu	Malaria	Leaf is smashed, mixed with water and decanted, then mixed with milk and one coffee cup is taken every morning.	Oral
				L	Li	Bloating	Leaves of Asparagus africanus& the root of Verbascum sinaiticum are pounded and mixed with water, decanted and dropped in to left nostril.	Nasal
Brassica carinata A.Br.	Brassicaceae	Gomenzer [A] Ija raafuu[O]	H	Se	Hu	Stomach problem	Crushed and eaten with Allium sativum by adding salt.	Oral
				Se	Hu	Cancer	The seed of Brassica carinata together with seed of Ricinus communis is crushed, powdered and mixed with honey and then creamed on affected part.	Dermal
Brassica nigra L.	Brassicaceae	Senafich[A] Sanaafica[O]	H	Se	Hu	Malaria	Powdered seed of Brassica nigra, chopped Allium sativum and Cicer arietinum seed [ Shimbura] are soaked with water and eaten after one day by enjera in the morning.	Oral
				Se	Hu	Intestinal parasites Amoeba	The semi-liquid condiment made of Brassica nigra called [awaze] is eaten with either injera or germinated bean seeds.	
Brucea antidysenterica Fresen.	Simaroubaeae	Waginos [Abalo[A] Qomanyoo [O]	Sh	L	Li	External parasites [lice]	Leaf of Brucea antidysenterica is pounded and mixed with water. The mixture is used to wash skin of cattle, donkey, mule and horse.	Dermal
				L	Hu	Wound and itches	Dried leaf of Brucea antidysenterica is pounded, mixed with butter and creamed the affected part until recovery.	Dermal
				R	Hu	Evil eye	Dried root of Brucea antidysenterica and Carissa spinarum are mixed together,	Nasal

							smoked & inhaled.	
				Se	Hu	Back pain	Seed of Brucea antidysenterica is boiled with milk and drunk for 3-4 consecutive days.	Oral
Buddleja polystachya Fresen.	Loganiaceae	Anfar[A] Adaaddii[O] Hanfaaree[O]	Sh	L	Li	Eye disease	Fresh Leaf of Buddleja polystachya is chewed and spitted on cattle eye.	Eye
				L	Hu	Wound	Fresh leaf of Buddleja Polystachya is pounded, powdered and applied on wound	Dermal
Calpurnia aurea [Ait.] Benth.	Fabaceae	Digita[A] Ceekaa [O]	Sh	L	Hu	Wound	Dried leaf of Calpurnia aurea is pounded, mixed with butter or honey and creamed on the wounded part.	Dermal
				L	Li	Scabies and Lice	Leaf of Calpurnia aurea, Croton macrostachyus and Justicia schimperiana are pounded, mixed with water and wash the body of the cattle every morning until the Scabies [itches] and parasites are eradicated.	Dermal
				L	Both	Snake bite	Leaf of Calpurnia aurea is squeezed and drop of the sap is given orally to cattle and to Human.	Oral
				L	Hu	Diarrhea	Fresh leaf of Calpurnia aurea is squeezed and the juice is drunk before breakfast.	Oral
Capsicum Annum L.	Solanaceae	Berbere [A] Qaaraa[O]	H	Fu/Se	Hu	Skin rash	Fruit and seed of Capsicum annumis pounded, powdered, mixed withbutter and creamed the infectedparts	Dermal
					Hu	Tonsillitis	Fruit and seed of Capsicum annumis pounded, powdered, mixed withoil, roasted and drunk	Oral
Carica papaya L.	Caricaceae	Papaya [A] Pappayya[O]	T	Fu	Hu	Heart problem	Fruit is eaten	Oral
				L	Hu	malaria	Leaf of Carica papaya and Allium sativum bulb are pounded together and made in the form of soup, boiled and mixed with honey and two cup are drunk.	Oral
				L	Hu	Intestinal parasites	Fresh leaves are boiled with water and cooled then drunk in the morning.	Oral
Carissa spinarum L.	Apocynaceae	Agam[A] Agamsa[O]	Sh	R	Hu	Evil eye	Root of Asparagus africanus, root, Lobelia rhynchopetalum bark, Artemisia abyssinica root and leaf, Allium sativum and Ruta chalepensis are crushed and smashed together and rolled by piece of cloth and tied on neck or arm.	Neck
				L	Hu	Head ache	Dried Leaf of Carissa spinarum is pounded and the smoke is used as treatment for head ache.	Nasal
				L	Hu	Stomach-ache	Leaf of Carissa spinarum is Pounded, mixed with honey. Two-three spoon is taken early in the morning before breakfast.	Oral

				L	Hu	Malaria	Fresh root is pounded, mixed with cold water, decanted and drunk after one day.	Oral
Citrus limon [L.] Burm.f.	Rutaceae	Lomi [A] Loomii [O]	Sh	L	Hu	Cough	The leaf of Citrus limon is pounded, powdered, mixed with milk, boiled and sugar is added. Then drink pure liquid during feeling of the pain.	Oral
				Fu	Hu	Athlete's foot	Fruit of Citrus limon is squeezed and creamed on affected part for continuous days.	Dermal
				Fu	Hu	Scabies[itches]	Fruit juice is squeezed and applied to the affected part.	Dermal
				Fu	Hu	Vomit	Fresh fruits are squeezed and the juice is drunk	Oral
Clematis simensis Fresen.	Ranunculaceae	Azo hareg[A] Hidda feetii[O]	Cl	L	Hu	Cancer	Fresh leaf is crushed, smashed, rolled by clean cloth and tied on hand.	Dermal
					Li	Horse itch	Leaf of Clematis simensis is squeezed and dropped on itched part.	Dermal
				R/L	Hu	Tonsillitis	Leaf of Clematis simensis is crushed, rolled in clean cloth and tied on neck.	Neck
				R/L	Hu	Wound	Dried root and leaf of Clematis simensis are pounded, powdered, mixed with butter and creamed on affected part.	Dermal
Clerodendrum myricoides [Hochst.] Vatke	Lamiaceae	Misirich[A] Maraasisaa[O]	Sh	R	Hu	Stomach-ache	Root of Clerodendrum myricoides, fruit of Croton macrostchys and root of Solanum adoense are crushed together and mixed with rotten butter and drunk.	Oral
				R	Hu	Headache	Root latex of Clerodendrum myricoides is dried, powdered, mixed with butter and creamed on head	Dermal
				R	Hu	Evil eye	Dried root is crushed and put on broken pot, and then fire is added to it and fumigated.	Nasal
				L	Hu	Eye problems	Leaves are pounded and powdered then sprayed to the eye.	Eye
Clutia abyssinica Jaub. & Spach	Euphorbiaceae	Fiyalefaj [A] Ulee loonii[O]	H	Se	Hu	Ring worm	Fresh seed is smashed, mixed with butter and creamed on affected part.	Dermal
				L	Hu	Ear disease	Leaf of Clutia abyssinica is pounded, squeezed and then its drop is added through the ear.	Ear
				R	Hu	Toothache	Fresh root is chewed and kept on teeth for some times.	Oral
Coffea Arabica L.	Rubiaceae	Buna[A] Buna[O]	Sh	Se	Hu	Diarrhea	Powder of roasted coffee bean is mixed with butter and eaten or drunk before breakfast for 3-4 days.	Oral
					Hu	Fire burn	Seed of Coffea arabica is roasted, pounded, powdered and applied on affected part.	Dermal
					Hu	Spider poison	Seed of Coffea arabica is Roasted, powdered, mixed with butter and painted.	Dermal
					Hu	Eye disease	Roasted seeds of Coffea Arabica is pounded together with Leaf of Trigonella foenum-graecum, mixed with butter and rubbed on the external eye.	Eye
Cordia africana Lam.	Boraginaceae	Wanza[A] Waddeessa[O]	T	B	Hu	Unstopped menstruation	Fresh bark is pounded, mixed with water, decanted and drunk with one coffee cup for three consecutive days	Oral
				L	Hu	Toothache	Fresh leaf is chopped, chewed with salt and the juice is swallowed.	Oral
				Fu	Hu	Intestinal parasites	The Fruit is eaten as food for the case of Intestinal parasites in the morning before food for 4-5 consecutive days.	Oral
Coriandrum sativum L.	Apiaceae	Denbelal[A] Dinbilaala[O]	H	Se	Hu	Cough	The seed together with butter and small quantity of water are boiled and drunk.	Oral
					Hu	Wound	Leaf of Coriandrum sativum is pounded with leaf of Croton macrostachys and	Dermal

							Rumex nervosus. Then creamed on wounded part for 2-3 days.	
Croton macrostychus Del.	Euphorbiaceae	Bisana [A] Bakkannisa [O]	T	Sht	Hu	Gonorrhea	Seven, nine or eleven shoot tips is cut, cooked together with Ruta chalepensis and one spoonful of the solution is drunk per a day for seven consecutive days.	Oral
				B	Li	Bloating	The bark of root is ground, mixed with water and given to the animal.	Oral
				L	Li	Scabies[itches]	Leaf of Croton macrostachyus with leaf of Brucea antidysenterica are crushed together and used as skin wash for calf.	Dermal
Cucumis ficifolius A.Rich.	Cucurbitaceae	Midirenbuay[A] Coolotoo[Hiddii hooloo][O]	Cl	R	Hu	Gonorrhea	Root of Cucumis ficifolius and root of Cyphostemma adenocaula are crushed and pounded together, mixed with water and decanted, then mixed with honey and drunk.	Oral
				R	Li	Cattle infection	Root of Cucumis ficifolius with leaf of Vernonia amygdalina are pounded together and mixed with cold water. Then given to cattle by tella material [merti].	Oral
				Fu	Hu	Stomach-ache	The root of Cucumis ficifolius is chewed and swallowed during the feeling of ach	Oral
				Sa	Hu	Ear pain	Sap of fruit is added to ear canal.	Ear
Cucurbita pepo L.	Cucurbitaceae	Duba [A]Dabaaqula [O]	Cl	Se	Hu	Tape worm	Seed of Cucurbita pepo is roasted and eaten before breakfast.	Oral
Cupressus lusitanica Mill.	Cupressaceae	YeFerenjtsid[A]Gaattiraa faranjii[O]	T	La	Hu	Ringworm	Latex of Cupressus lusitanica is creamed on affected part of body.	Dermal
Cynoglossum coeruleum [Hochst.ex.A.rich.]	Boraginaceae	Chigogit[A]Maxxannee [O]	H	L	Hu	Headache	Fresh leaf of Cynoglossum coeruleum is massaged and sniffed.	Nasal
				R		Body swelling	Fresh root of Cynoglossum coeruleum is chewed and sprayed on swollen part	Dermal
				L		Scabies[Itches]	Juice of fresh leaves is applied on the skin.	Dermal
Cyphostemma adenocaula [Steud. ex A. Rich.] Des. ex Wild & Drum.	Vitaceae	Aserkush[A]Hidda Bofaa[O]	Cl	Tu	Hu	starvation	Tuber is cooked and eaten during famine	Oral
				Ag	Li	Infection on Neck[livestock]	Tied on the livestock's neck.	Neck
Datura stramonium L.	Solanaceae	Astanagir[A]Manjii [O]	H	L	Hu	Dandruff	Leaf is crushed with leaf of Myrtus communis & the powder is rubbed over the head after having cut the hair.	Dermal
				R	Hu	Headache	Roots of Datura stramonium is pounded with leaf of Ocimum gratissimum and sniffed Nasally.	Nasal
				Fu	Hu	Scabies[itches]	About 2-3 spoons of Powdered fruit of Datura stramonium is mixed with butter and creamed.	Dermal

				L	Hu	Wound	Fresh leaf of Datura stramonium is squeezed and creamed the affected [wounded] part of the body.	Dermal
Discopodium penninervum Hochst.	Solanaceae	Almit[a]	Sh	L	Li	Leech	The leaf of Discopodium penninervum is pounded, mixed with little water and the decanted is given to cattle.	Oral
					Li	Bloating	The fresh leaf of Discopodium penninervum is pounded, squeezed, mixed with water, decanted and given to cattle.	Oral
Dodonaea angustifolia L.	Sapindaceae	Kitkita[A] Ittacha[O]	Sh	L	Hu	Diarrhea	Fresh leaves are pounded, mixed with water, decanted and drunk.	Oral
					Hu	Skin rash	Fresh leaves are crushed, mixed with water and painted the skin.	Dermal
					Li	Wound	Dried leaves of Dodonaea angustifolia are powdered and sprayed on the wound of pack [domesticated] animals.	Dermal
						Ecto-parasities	Fresh leaves are crushed and pounded with water, then leaf extract is given Orally and pounded leaves are painted on the cattle body.	Dermal
Dombeya torrida [J. F. Gmel.] P. Bamps	Sterculiaceae	Wulkifa [A] Daannisa[O]	T	L	Hu	Fire burn	Fresh leaf of Dombeya torrida is squeezed and creamed on affected part.	Dermal
						Abdominal pain	Dried leaf powder is mixed with honey and given Orally.	Oral
Dovyalis abyssinica [A.Rich.]Warb.	Flacourtiaceae	Koshim[A] Koshommii [O]	Sh	Fu	Hu	Intestinal Parasites	Its fruit is eaten as food for the case of intestinal parasite before breakfast every morning	Oral
Echinops kebericho Mesfin	Asteraceae	Kerebicho [A] Qarabichoo [O]	H	R	Hu	Evil eye	Root concoction together with Leaf of Artemisia abyssinica is added to a burning fire and smoked to the patient.	Nasal
				R	Hu	Rabies	Root concoction together with root of Ricinus communis is boiled and decoction is drunk.	Oral
				R	Hu	Evil Spirit	Root is Crushed, heated on fire and its smoke is inhaled.	Nasal
				R	Hu	Tetanus	Root of E. kebericho with bark of Croton macrostachys is pounded, powdered together, mixed with honey and then one cup of the mixture is taken for 3 days.	Oral
Embelia schimperi Vatke	Myrsinaceae	Enkoko [A] Haanqu [O]	Sh	Se	Hu	Tape worm	Seed of Embelia schimperi is dried and powdered, mixed with water, two glasses are taken in the morning before food and stay for 6-7 hours.	Oral
				Fu	Hu	Epilepsy	Fruit of Embelia schimperi with seed of Guizotia abyssinica is crushed, powdered, mixed with local alcohol "tej" and drunk	Oral
Erica arborea L.	Ericaceae	Asta[A] Maxaxee [O]	T	Sht	Hu	Giardiasis	Shoot is boiled and the decoction is drunk for 3-4 days consecutively.	Oral
				L	Hu	Wound	The powder of dried leaves is mixed with butter & rubbed on the affected part.	Dermal
Eucalyptus globulus Labill.	Myrtaceae	Nech bahirzaf[A] Bargamoo	T	L	Hu	Influenza	The Leaf of Eucalyptus globulus is chopped and Boiled with water and inhale the vapor repeatedly.	Oral/Nasal

		adii[O]		St	Hu	Fibril illness	The young leaves and immature stem tip are boiled in water and the steam is inhaled by the patient during the sleep time until recovery.	Nasal
				L	Hu	Malaria	Dried leaf is put on fire and smoked	Nasal
				L	Hu	Cough	Leaf is boiled with sugar and one tea cup of syrup is drunk.	Oral
Euphorbia abyssinica. Gmel	Euphorbiaceae	Kulkual [A] Adaammii [A]	T	Fw	Hu	Wound	Fresh flower of Euphorbia abyssinica is squeezed and creamed on affected part	Dermal
				B	Hu	Ascariasis	Fine powder of pounded bark of Euphorbia abyssinica and Croton macrostachyus is mixed with water, decanted and taken at meal time.	Oral
				La	Hu	Gonorrhoea	Very small amount of the milky latex is mixed with red teff flour, baked and eaten for three consecutive days.	Oral
Guizotia abyssinica L.	Asteraceae	Nug[A] Nuugii[O]	H	Se	Hu	Kidney Problem	Seed of Guizotia abyssinica is roasted, pounded and mixed with powdered leaves of Thymus schimperii and boiled. Then one coffee cup is drunk for 3-5 days.	Oral
				Se	Hu	Swelling	Seed is roasted, pounded and powdered, then boiled with salt and the syrup is drunk.	Oral
				Se	Li	Leech	More than one litter powdered seed mixed with water is given to cattles for three days continuously.	Oral
Hagenia abyssinica [Bruce] J. F. Gmel.	Rosaceae	Koso [A] Heexoo[O]	T	Fu	Hu	stomach-ache	Fresh fruit is crushed, squeezed and the juice is mixed with tella and drunk.	Oral
				Se	Hu	Venereal disease [abalazar]	Dried Seed is powdered, mixed with tela and drunk.	Oral
				Se	Hu	Eczema	Calf horn and roasted Hagenia abyssinica seed are pounded together, mixed with butter and creamed on affected body part.	Dermal
				Se	Hu	Tape worm	The seed of Hagenia abyssinica is crushed, powdered, mixed with milk, boiled and drunk for five days before breakfast.	Oral
Hypericum revolutum Vahl	Hypericaceae	Amja [A] Muka Foonii[O]	Sh	L	Hu	Erythroblasts	Leaf is boiled with water and the decoction is drunk.	Oral
				Sht	Hu	Dandruff	Shoot is crushed, squeezed and the sap is creamed after having cut hair.	Dermal
				L/R	Li	Rabies	The root and leaf of Hypericum revolutum together with roots of Rumex nervosus, Phytolacca dodecandra, Brucea antidysenterica, leaf and bark of Croton macrostachyus are pounded together, mixed with water and given to cattle.	Oral

Jasminum abyssinicum L.	Oleaceae	Tembelel Biluu[O]	CI	L	Hu	Eye disease	The leaves of Jusminium abysinicum and seven pieces of immature stems of Olea europaea are ground and powdered together, mixed with water and drop of mixture is applied on infected part for 3 days.	Eye
Juniperus procera Endl.	Cupressaceae	Ye abesha tsid[A] Gaattiraa[O]	T	B	Hu	Tonsillitis	Fresh stem bark is chewed and the juice is swallowed.	Oral
				B	Hu	Malaria	Dried bark is boiled with water and the decoction is drunk.	Oral
				L	Hu	Pneumonia	Fresh leaf is Crushed and steeped in cold water and the infusion is drunk.	Oral
Justicia schimperiana [Hochst. exNees] T. Anders	Acanthaceae	Simiza[sens el][A] DHummuugaa[O]	Sh	L	Li	Hen disease [Cocoidiosi]	The leaf [shoot] is crushed, mixed with water and decanted; 'Enjera' [bread] is soaked with the solution and given to the hen.	Oral
				L/R	Li	Blackleg	Leaf and root of Justica schimperiana is pounded with dried fruit of Ricinus communis and the solution is given to cattle.	Oral
				L	Hu	Leg swelling	Fresh Leaf is heated on fire; salt is added to it and tied on the swollen part.	Dermal
				L	Li	Lice	Leaf of Justicia schimperiana with leaf of Calpurina aurea are used to wash the body of cattle to remove the lice.	Dermal
Kalanchoe petitiiana A. Rich.	Euphorbiaceae	Endawula [A] Bosoqqee [O]	H	R	Hu	Bone fracture	Fresh leaf is squeezed; the juice is mixed with butter and applied topically on fractured part.	Dermal
				L	Hu	Wound	Fresh leaf is heated with fire, squeezed and the juice is dropped on wounded part	Dermal
				R	Hu	Ear disease	Fresh root of Kalanchoe petitiiana is squeezed and few drops are added through ear canal.	Ear
Kosteletzkya begonifolia Ulbr.	Malvaceae	Yemegerem [A]	H	R	Li	For anthrax	Fresh leaf of Kosteletzka begonifolia is pounded squeezed mixed with "tella" and drunk.	Oral
				R	Hu	For body swelling	Fresh leaf of Kosteletzka begonifolia is pounded squeezed mixed with "tella" and drunk.	
Linum usitatissimum L.	Lineaceae	Telba[A] Talbaa[O]	H	Se	Hu	Wound	Seed of Linum usitatissimum is pounded, mixed with honey and creamed on wounded part	Dermal
				Se	Li	Retained placenta	Seed of Linum usitatissimum is powdered and half of a glass of the powder is dissolved in water and given to cattle.	Oral
				Se	Hu	Amoeba	The pounded seed is mixed with water and drunk before breakfast.	Oral
				Se	Hu	Gastritis	The seed of Linum usitatissimum is pounded, mixed water and sugar and then drunk during feeling the pain.	Oral
Lobelia rynchopetalum Hemsl.	Lobeliaceae	Jibira[A]	Sh	B and	Hu	Evil eye	The bark and root of Lobelia rynchopetalum is crushed , mixed with little water and sniffed at the sickness time or the dried root is tied with piece of cloth	Nasal and

				R			around the neck or put it in the pocket.	Dermal
				R	Hu	Scabies[itches]	Root of Lobelia rynchopetalum is dried, powdered, mixed with butter, rub the body for some days and finally taking a shower at the end of the day.	Dermal
Lepidium sativum L.	Brassicaceae	Feto[A] Feecoo[O]	H	Se	Hu	Bloating	Seed of Lepidium sativum and bulb of Allium sativum are pounded together, mixed with water and given to cattle.	Oral
				Se	Hu	Dysentery	The seed of Lepidium sativum is ground, mixed with milk, then filtrate and the solution is drunk.	Oral
				Se	Hu	Malaria	Seed of Lepidium sativum, bulb of Allium sativum and rhizome of Zingiber officinale are pounded together and given to Human with honey.	Oral
				Se	Hu	Tonsilitis	Seed of Lepidium sativum and bulb of Allium sativum are pounded together and given to Human with honey.	Oral
				Se	Hu	Fibril illness	Dried seeds are powdered and added in to fire and smoked to the patient.	Nasal /Oral
				St	Hu	Hemorrhoids	Stem is heated on fire and used to burn the affected part	Dermal
Maesa lanceolata Forssk.	Myrsinaceae	Shwaria[ kelawa ] [A] Abbayii [O]	Sh	L and Se	Li	For leeches	Dried leaf and fruit of Maesa lanceolata are crushed, pounded and the powder is added to water to kill leeches	Oral
				R	Li	Retention of placenta	Root of Maesa lanceolata is boiled with seed of Linum usitatissimum and given to cattle.	Oral
				L	Hu	Eczema[chife]	Dried leaf powder together with the powder of leaf of Croton macrosrachyus are pasted with butter and applied for seven days consecutively.	Dermal
Melia azedarach Forssk.	Meliaceae	Nim[A] Niimii[O]	T	L	Hu	Malaria	Chewing and swallowing the juice of fresh leaf.	Oral
				B	Li	Anthrax	The fine powder of dried bark is added to a glass of water and applied through the mouth.	Oral
				Sht	Hu	Tooth ache	Young shoot tip is chewed and kept on the teeth.	Oral
Myrtus communis L.	Myrtaceae	Ades[A] Adasii[O]	Sh	L	Both	Intestinal parasites	Dried leaf of Myrtus communis is pounded, added to tella and drunk to expel intestinal parasites.	Oral
				L	Hu	Dandruff	The leaf of Myrtus communis is powdered, mixed with butter and creamed on head after is cut.	Dermal
				L	Hu	Scabies[itches]	Dried powder is mixed with butter & applied on the affected part	Dermal
Nicotiana tabacum L.	Solanaceae.	Timbaho[A] Tamboo[O]	H	L	Hu	Snake bite	Leaf of Nicotiana tabacum is crushed and squeezed, mixed with water and the Juice is drunk to expel the poison by vomiting.	Oral
				L	Li	Bloating	Leaf and root of Nicotiana tabacum is dried, powdered, mixed with salt, water and made as bread. Slice is given to cattle before they drink water for three days.	Oral



				St/ L	Li	Leech	The young stems and or leaf is ground, add salt, and then one glass of the mixture is given every morning for three days orally or through the nose.	Oral
				L	Li	Internal parasites	Leaf of Nicotiana tabacum is pounded with root of Carissa spinarum and mixed with water. One merit solution is given to calf.	Oral
Nigella sativa L.	Ranunculaceae	Tikur azmud[A] Abasuuda gurraacha [O]	H	Se	Hu	Depression	Seed is added to tea and drunk to stimulate mental.	Oral
				Se	Hu	Tonsillitis	Seed of Nigella sativa is pounded, powdered and added to coffee. Then drunk for 3-4 consecutive days.	Oral
				Se	Hu	Stomach-ache	Seed is crushed and pounded with Ruta chalepensis, Lepidium sativum and Allium sativum then mixed with honey and a spoonful of the mixture is taken before breakfast.	Oral
Ocimum basilicum L.	Lamiaceae	Zikakibe[A] Gosobila[O]	H	L	Hu	Headache	Leaf of Ocimum basilicum is crushed and sniffed.	Nasal
				L	Hu	Malaria	Leaf of Ocimum basilicum and bulb of Allium sativum are pounded together and eaten with honey in the morning.	Oral
				L	Li	Bloating	Its fresh leaves with the bulb of Allium sativum and salt are ground together and then one litter of the solution is given to cattle.	Oral
Ocimum lamiifolium Hochst. ex Benth.	Lamiaceae	Damakese[A] Damakasee[O]	Sh	L	Hu	Febrile illness	Fresh leaf of Ocimum lamiifolium together with leaf of Eucalyptus globulus, is pounded, mixed with water and drunk or the patient can inhale the vapor of the boiled mixture.	Oral
				L	Hu	Headache	Leaf of Ocimum lamiifolium is massaged and sniffed.	Oral
Olea europaea L. subsp. cuspidata [Wall. ex G. Don] Cif.	Oleaceae	Weyra[A] Ejersa[O]	T	L	Both	Eye disease	Leaf is Crushed and squeezed. Then the sap is dropped by piece of clean cloth on eye.	Eye
				St	Hu	Wound	Fresh stem is heated on fire and the oily liquid produced from the stem is applied on the wound.	Dermal
				St	Hu	Gastiritis	Oily liquid produced from the stem is drunk after meal for three consecutive days.	Oral
				B	Hu	Haemorrhoid	The bark is heated on fire and held on the pain area.	Dermal

Opuntia ficus-indica [L.] Miller	Cactaceae	Balas [Ashewa kulkual[A] Adaammii [O]	Sh	L	Hu	Ear disease	Leaf of Opuntia ficus-indica is squeezed and the sap is dropped through ear canal safely.	Ear
				Fu	Hu	Stomach-ache	Ripened fruit is eaten	Oral
Ostostegia integrifolia Benth.	Lamiaceae	Tunjit[A] Xunjiitii[O]	Sh	L	Hu	Stomach-ache	Leaf of Ostostegia integrifolia and leaf of Solanum adoense are crushed, squeezed together and the juice or sap is drunk.	Oral
				Hu		Fibril illness	Dried leaf is added to fire and fumigated	Oral and Nasal
				Hu		Tonsillitis	Fresh leaf of Ostostegia integrifolia is squeezed and half of coffee cup is drunk	Oral
Osyris quadripartita Decn	Santalaceae	Keret [A] Waatoo [O]	Sh	St	Hu	Stomach-ache	Fresh stem bark of is chewed & the solution is swallowed.	Oral
				L	Hu	Wound	Dried or fresh leaf of is crushed, powdered and then applied on the wound part	Dermal
Phytolacca dodecandra L' Herit.	Phytolaccaceae	Endod[A] Handoodee[O]	Sh	Se	Hu	To stop pregnancy	Seed of Phytolacca dodecandra are crushed, smashed, squeezed and the juice is drunk.	Oral
						Malaria	Fresh root is smashed, mixed with water, decanted and drunk in the morning.	Oral
				R	Li	Bilharziha	Fresh root is chewed and swallowed.	Oral
						Stomach-ache	Fresh root is crushed and pounded with water, then filtered and drunk for four days	Oral
Pittosporum viridiflorum Sims	Pittosporaceae	Tebera[A]Dambii[O]	T	L	Hu	Dandruff	Leaf is crushed, smashed and mixed with butter and creamed after having cut hair.	Dermal
Plantago lanceolata L.	Plantaginaceae	Gortab Ye wusha milas Wenberet[A] Qorxobbii [O]	H	L	Hu	Wound	Fresh leaf of Plantago lanceolata and bulb of Allium sativum are crushed, smashed together and rubbed on wounded part.	Dermal
						Fibril illness	Rub the body with the squeezed leaves	Dermal
						Skin cut	Fresh leaf of Plantago lanceolata is smashed, squeezed and three to four drops of the exudate is added to skin cut.	Dermal
Prunus persica [L.] Batsch	Rosaceae	Kok[A] Kookii [O]	T	L	Hu	Constipation	Fresh leaf of Prunus persica is crushed and mixed with water and given orally for continuous days until it gets relieve.	Oral
				L	Hu	Diarrhea	Leaf is pounded, mixed with water and decanted, and then one cup of local tella is given to Human.	Oral
				Ep	Hu	For delayed	Leaf of epiphyte is cut by her Husband. Then crushed, mixed with water,	

						pregnancyWoman[Infertile]	decanted, drunk at bed time and continuing sexual intercourse.	Oral
Pterolobium stellatum[Forsk]Brenan	Fabaceae	Kentafa[A] Harangamaa [O]	Sh	L	Hu	Goiter	Crush the leaves and mix with butter. Then apply the paste and tie it on the neck.	Dermal
				L	Hu	Evil eye	Leaf of Pterolobium stellatem and Ruta chalepensis are pounded, mixed with water and one coffee cup is drunk.	Oral
				R	Hu	Sudden sickness	Root of Pterolobium stellatem is dried, powdered and preserved. One spoon of the powder is mixed with alcohol and given to human.	Oral
Rhamnus prinoides L. Herit.	Rhamnaceae	Gesho[A] Geeshoo[O]	Sh	L	Hu	Teeth-ache	Leaf is chewed and kept on the teeth.	Oral
				L	Li	Leech	Leaf is crushed, dried and soaked with the leaves of a Solanumtuberosum at least for 2 days & then given small droplets through left nostril & then much more through their mouth.	Oral and Nasal
				L	Hu	Tonsillitis	Young leaves are chewed and the juice is swallowed.	Oral
Ricinus communis L.	Euphorbiaceae	Gulo[A] Qobboo[O]	Sh	Se	Hu	Headache	Seed is pounded, mixed with butter, and tied on head.	Dermal
				Se	Li	Anthrax	Fresh Fruit is pounded, mixed with water and given for cattle to drink.	Oral
				Se	Hu	Impotency	Seeds are pounded, mixed with small quantity of Aloe spp. latex and drink one coffee cup before bed time for 3-4 days.	Oral
				L	Li	Bloating	Fresh leaf is pounded, mixed with water; salt is added and given to cattle Orally.	Oral
Rosa abyssinica Lindley	Rosaceae	Kega[A] Goraa[O]	Sh	L	Hu	Ascariasis	Fresh leaf is pounded, mixed with water and a cup of the mixture is drunk.	Oral
				Fu		Hypertension	Fresh fruit with the leaves of Otostegia integrifolia are ground, powdered, mixed with water and one coffee cup is drunk during the sick time.	Oral
				Fw/L		Erythroblasts	The flower and leaf of Rosa abyssinica is eaten for some days.	Oral
Rumex abyssinicus Jacq.	Polygonaceae	Makmako[A] Dhangaggoo Fardaa[O]	H	R	Hu	“Ayne tila”	Root of Rumex abyssinicus and root of Verbascum sinaiticum are crushed together and mixed with honey. Then drunk before breakfast for three consecutive days.	Oral
				R	Hu	Hypertension	Dried root of Rumex abyssinicus is pounded , added to tea and drunk	Oral
				R	Hu	Ring worm	The root of Rumex abyssinicus with root of Rumex nepalensis is pounded, powdered, mixed with solution of Citrus limon and creamed on affected part.	Dermal
Rumex nepalensis Spreng.	Polygonaceae	Lut [A] Tultii[O]	H	R	Li	Retained placenta	Latex of Rumex nepalensis root is washed out by water and given to animal [e.g. sheep, cow]	Oral

				R	Hu	Stomach-ache	Root is Chewed and the sap is swallowed.	Oral
				R	Hu	Wound	Root is chopped, dried, powdered and applied on wounded part.	Dermal
				L	Hu	Spider poison	Leaf of Rumex nepalensis is directly rubbed on affected skin.	Dermal
				R	Hu	Internal Parasite	The root is Chewed and swallowed or boiled in the water and one glass of the solution is drunk.	Oral
				R	Hu	Body swelling	Dry/fresh root is chewed and put on the swollen part.	Oral
Rumexnervosus Vahl	Polygonaceae	Embacho [A] Dhangaggoo	Sh	L	Hu	Wound	Dried root bark is crushed, pounded, mixed with butter and placed on the wound.	Dermal
				R		Skin rash	Root of Rumex nervosus is dried and powdered. One coffee cup of powder is mixed with butter and creamed on affected skin	Dermal
				L	Hu	Snake bite	The leaf of Rumex nervosus is chewed and the solution is swallowed during bite time.	Oral
				L/ St	Li	Lice	Rumex nervosus together with Sida schimperiana is used to wash the calf to remove external parasites and Scabies [itches].	Dermal
Ruta chalepensis L.	Rutaceae	Tenadam[A] Cilaattama[O]	H	L	Hu	Stomach-ache	Fresh leaf of Ruta chalepensis mixed with sugar & powder of Vicia faba, then boiled and drunk [or chewing and taking the sap].	Oral
				B/L	Li	Cocoidiosis [Bilii]	Bark and leaf of Ruta chalepensis and root of Justicia schimperiana are pounded together and given to hen with injera.	Oral
				Se	Hu	Evil eye	The seed of Ruta chalepensis with Allium sativum are finely crushed together and sniffed at the sickness time	Nasal
Salix macronata THunb.	Salicaceae	Lahaya[A] Alaltuu[O]	T	Sht	Hu	Hemorrhage	Fresh shoot is crushed, squeezed and the juice is rubbed on the affected part.	Oral
				L	Hu	Wound	Fresh leaf of Salix macronata is pounded and placed on wounded part	Dermal
				L	Li	Joint dislocation	The leaf is ground along with young stem, mixed with bread and given to the cattle in problem	Oral
Schinus molle L.	Anacardiaceae	Kundoberbere[A] Qundobarbarree[O]	T	L/Fu	Li	Eye disease	Fesh Leaf and fruit of Schinusmolle are chewed and spitted on cattle, equines, goat and sheep eye.	Eye
				Se	Hu	Tonsillitis	Dried seed of Schinusmolle is pounded, powdered, mixed with honey and then drunk	Oral
Sesamum indicum L.	Pedaliaceae	Selit[A] Saliixa[O]	H	Se	Hu	Ear defect	Extract oil from the seed and drop the extract in canal the ear.	Ear
Sidas chimperiana Hochst. Ex A. Rich.	Malvaceae	Cifrig[A] Harmellaa[O]	H	R	Hu	Syphilis [ketegn]	Fresh Root of Sidaschimperiana and root of Solanumadoense are crushed together and mixed with water and decanted. Then mixed with honey and drunk.	Oral
				R	Li	Horse disease	Ground, mixed with water and drunk or applied through the nostrils.	Nasal

				R	Hu	Evil eye	Root is fumigated every evening to patient until recovery.	Nasal
Solanum adoense [Hochst] ex A. Rich.	Solanaceae	Zerch Enbuay[A] Hiddii baddaa[O]	Sh	Fu	Hu	Wound	Adding the drop of Solanum adoense fruit juice on wounded part	Dermal
				Fu	Li	Bloat	Fruits of Solanumadoense are Squeezed on cattle feed.	Oral
				L	Li	Anthrax	Dried leaf of Solanum adoense is crushed, pounded, mixed with tella and given to cattle.	Oral
Solanum incanum L.	Solanaceae	Enbuay[A] Hiddii loonii[O]	Sh	R	Hu	Snake bite	Dried root powder is drunk with coffee.	Oral
				Sa	Hu	Infection caused by spine in leg	The sap of fresh fruit is squeezed & dropped on affected part.	Dermal
				L	Hu	Nasal bleeding	The leaf of Solanum incanum is ground, powdered and sniffed several times until recovery.	Nasal
				L	Li	Urination problem	The leaf of Solanum incanum and that of Cuminum cyminum are smashed together, mixed with water and given for horse, donkey, and mules to drink	Oral
Solanecio gigas [Vatke.]C. Jeffery	Astraceae	Yeshikoko gomen[Mog ne qitel[A]] Jirma Jaldeessaa[O]	Sh	L	Li	Bloating	The leaf of Solanecio gigas is pounded, squeezed and the juice is added through the nose	Nasal
				L	Li	Lice	Leaf of Solanecio gigas is used to wash hair of calf as lice killer.	Dermal
Stephania abyssinica [Dillo&A.Rich.]Walp.	Menispermaceae	Ye ayit hareg[Chewchawit [Etse Iyesus[A] Hidda kalaalaa[O]]	Cl	L	Hu	Wound	Leaf of Stephania abyssinica is pounded and a small amount is added to wound.	Dermal
				R	Li	Rabies	Dry root of Stephaniaabyssinica is powdered and baked with teff flour and given to cattle.	Oral
				L	Li	Pasturolosis	Dried root and leaf of Stephaniaabyssinica is powdered together, mixed with water and given to the animal [cattle, goat or sheep].	Oral
Syzgium aromaticum L.	myrtaceae	Kirunfud[A] Qurunfudii[O]	T	Fu	Hu	Impotency	Dried Syzgium aromaticum is crushed, mixed with goat milk and boiled. Then the decoction is drunk.	Oral
				Fu	Hu	Vomit	Fruit of Syzgiumaromaticum with rhizome of Zingiberofficinale is boiled and drunk.	Oral
Tanacetum cinerariifolium [Trev]. Sch. Bip.	Asteraceae	Kaba[A]	H	L	Hu	Fibril illness	The leaf of Tanacetum cineratiifolium is squeezed and drunk	Oral
Trigonellafoenum-Graecum L.	Fabaceae	Abish[A] Sunqoo[O]	H	Se	Hu	Leg wound	Seed of TrigonellafoenumGraecum, seed of Faciafaba and seed of Linumusitatissimum are finely ground together, rolled in piece of cloth and tied on leg	Dermal

				Se	Hu	Body swelling	The seed of Trigonellafoenum-graecum is crushed, powdered, mixed with honey and little water, then boiled like “porage” and eaten	Oral
				Se	Hu	Bone fracture	Seed is powdered; water is added to flour to make the paste, and then applied to the broken bone.	Oral
Thymus schimperi Ronniger	Lamiaceae	Tosign[A] Xasee[O]	H	Ag	Hu	Cough	Its leaves, root and bark are ground together, powdered and mixed with water, one coffee cup [sini] is drunk during pain time.	Oral
				L	Hu	Stomach-ache	Leaf is boiled with leaves of Foeniculumvulgare and one tea cup is taken.	Ora
				L	Hu	Hypertension	Leaf is boiled with sugar and drunk.	Oral
Urera hypselodendron [A. Rich.] Wedd.	Urticaceae	Lankuso[A] Laanqisaa dhogonu [O]	Cl	B	Li	Anthrax	Bark/shoot of Urera hypselodendron is crushed, smashed, mixed with powder of Trigonella foenum-graecum or with egg and given to cattle.	Oral
Urtica simensis Steudel	Urticaceae	Sama[A] Doobbii [O]	H	L	Hu	Gastritis, Heart Disease	Eat in the form of stew [‘wot’] against gastritis & heart disease.	Oral
				R/L	Hu	Gonorrhea	The root and leaves of Urticasimensis with the bark of Crotonmacrostachyus are pounded, powdered, mixed with little water ,filtered, then a cup of filtrate is drunk for 5 days in the morning	Oral
Verbena officinalis L.	Verbenaceae	Atuch[A] Derguu[O]	H	L	Hu	For tonsillitis	Leaf of Verbenaofficinalis is Crushed, smashed, mixed with butter and creamed around neck.	Dermal
				R	Hu	Fibril illness	Root of Verbenaofficinalis, together with the root of Carissaspinarum and root of Rutachalepensis, are fumigated to the patient.	Nasal
				R	Hu	Diarrhea	Root of V. officinalis and root of Phytolaccadodecandra, bark of Crotonmacrostachyus are pounded, mixed with water, decanted and drunk after a day.	Dermal
Verbascum sinaiticum Benth.	Scrophulariaceae	Kutina[ Ye Ahya joro[A] Gurra Harree [O]	H	R	Li	Horse disease	Fresh Root of Verbascumsinaiticum is collected from three places, smashed, mixed with water, decanted and given to horse	Oral
				L	Hu	Impotency	Chopped Leaf of Verbascum sinaiticum is rolled by clean piece of cloth and tied around male sex organ to erect it.	Dermal
				R	Hu	Wound	The root of Verbascum sinaiticum is crushed, powdered, mixed with butter and creamed on affected part.	Dermal
Vernonia amygdalina Del.	Asteraceae	Gerawa[A] Eebicha [O]	T	L	Hu	Malaria	Crushed leaves of Vernonia amygdalina concocted with leaves of Ruta chalepensis. One cup is served as a drink for 3-5 days with cold water in the morning.	Oral
				L	Hu	Skin infection	The leaf of Vernonia amygdalina is pounded and the patient body is washed by the plant or the leaf of is used as a soap to wash the infected body.	Dermal
				L	Li	Bloating Urine retention	Fresh leaves is squeezed, mixed with water, and then given to cattle in morning and at night until recovery.	Oral

Vernonia hymenolepis A.Rich.	Asteraceae	Weynagift [A] Sooyama[O]	Sh	L	Hu	Gonorrhea	Leaf twig of Vernonia hymenolepis and bark of Croton macrostachyus are pounded together, mixed with honey and 1-3 spoons is taken in the morning before breakfast.	Oral
				L	Hu	Wound	Leaf of Vernonia hymenolepis is squeezed and the sap is dropped on the wound.	Dermal
Zehneria scabra [Linn.f.] Sond.	Cucurbitaceae	Haregresia[Etse sabeq] Nech Hareg[A] Qorii Sinbiraa[O]	Cl	L	Hu	Swelling	Leaf and bark of Zehneria scabra and leaf of Rumex nervosues are pounded together, rolled in clean cloth, and tied on swelling.	Dermal
				L	Hu	Fibril illness	The leaf of Zehneria scabra is pounded, squeezed and mixed with sugar and drunk one cup /or the stem is boiled and inhaled	Oral
				R	Hu	Sudden sicknes	Fresh root is pounded, mixed with water and drunk.	Oral
				L	Hu	Dandruff	Fresh leaf is squeezed to make juice and creamed after hair is cut.	Dermal
Ziziphus spina- christi [L.] Desf.	Rhamnaceae	Gaba[A]	Sh	L	Hu	Dandruff	The leaf of Zizlphus spina-christi is pounded, powdered, mixed with butter and then creamed.	Dermal
Zingiber officinale Roscoe	Zingiberaceae	Jinjible[A] Zinjibila[O]	H	Rh	Hu	Malaria	Rhizome of Zingiber officinale and bulb of Allium sativum are pounded together and eaten with honey.	Oral
					Hu	Stomach ache	Rhizome of Zingiber officinale together with Leaf of Vernonia amygdalina and bulb of Allium sativum are pounded and eaten with honey.	Oral

**Appendix 3:** List of the medicinal plant families and corresponding number of genera and species in the study area

**Table 15**

No.	Family	No. of genera	No. of plant species	No	Family	No. of genera	No. of plant species
1.	Acanthaceae	1	1	27	Menispermaceae	1	1
2.	Alliaceae	1	2	28	Myrsinaceae	2	2
3.	Aloaceae	1	1	29	Myrtaceae	3	3
4.	Amaranthaceae	1	1	30	Oleaceae	2	2
5.	Anacardiaceae	1	1	31	Pedaliaceae	1	1
6.	Apiaceae	1	1	32	Phytolaccaceae	1	1
7.	Apocynaceae	1	1	33	Pittosporaceae	1	1
8.	Asparagaceae	1	1	34	Plantaginaceae	1	1
9.	Asteraceae	6	7	35	Polygonaceae	1	3
10.	Boraginaceae	2	2	36	Ranunculaceae	2	2
11.	Brassicaceae	2	3	37	Rhamnaceae	2	2
12.	Cactaceae	1	1	38	Rosaceae	3	3
13.	Caricaceae	1	1	39	Rubiaceae	1	1
14.	Cucurbitaceae	3	3	40	Rutaceae	2	2
15.	Cupressaceae	2	2	41	Salicaceae	1	1
16.	Ericaceae	1	1	42	Santalaceae	1	1
17.	Euphorbiaceae	5	5	43	Sapindaceae	1	1
18.	Fabaceae	4	4	44	Scrophulariaceae	1	1



					e		
19.	Flacourtiaceae	1	1	45	Simaroubaceae	1	1
20.	Hypericaceae	1	1	46	Solanaceae	5	6
21.	Lamiaceae	4	5	47	Sterculiaceae	1	1
22.	Lineaceae	1	1	48	Urticaceae	2	2
23.	Lobeliaceae	1	1	49	Verbenaceae	1	1
24.	Loganiaceae	1	1	50	Vitaceae	1	1
25.	Malvaceae	2	2	51	Zingiberaceae	1	1
26.	Meliaceae	1	1				

**Appendix 4:** List of human diseases in the study area

**Table 16**

No	Disease treated	Total of plant species used	Total percentage [%]
1.	Wound	25	26.88
2.	Malaria	14	15.05
3.	Stomach-ache	13	13.97
4.	Body swelling and Evil eye	10	10.75
5.	Tonsillitis	9	9.68
6.	Fibril illness, Scabies[itche] and Skin rash	7	7.52
7.	Cough, dandruff, diarrhea and headache	6	6.45

8.	Eye disease ,gonorrhoea ,intestinal parasites and tape worm	5	5.38
9.	Snake bite	4	4.30
10.	Amoeba,bone fracture, fire burn, hypertension, impotency, rabies,skin cut, sudden sickness andtoothache	3	3.22
11.	Asthma, cancer ,heart disease, hemorrhoids ,nasal bleeding, pneumonia, ring worms ,urination problem and vomit	2	2.15
12.	Abortion[ To stop pregnancy],back pain, bilharzia, ear defect, goiter,infertility, retained placenta and syphilis	1	1.07

**Appendix 5:** List of livestock diseases in the study area

**Table 17**

No.	Disease treated	Local name[Afaan oromoo]	No. of plant species used	Percent [%]
1	Bloating	Dhitahuu garraa	10	10.75
2	Anthrax and Leech	Abasanga Alekit	6	6.45
4	Ectoparasite [lice]	Ye wech tegegna	5	5.38
5	Rabies	Dhukkubaa sarre	3	3.22
6	Erythroblasts, horse disease, retained placenta , coccidiosis	Machangaf ,Dhukkuba fardaa,yengdelej sizegey and Dhukubaa indaqoo	2	2.15
7	Blackleg ,horse itch,pasturolosis and eye disease	Abagorba, Epizootic lymphagities,Gororsa, Aynebesheta	1	1.07

**Appendix 6:** Some of the cultivated and wild grown medicinal plants widely traded in the market for different uses in addition to medicinal values in Wonch district.

**Table 18**

<b>No</b>	<b>Scientific Name of the medicinal plant</b>	<b>Local name[Afan oromo]</b>	<b>Used for</b>
1.	Allium cepa	Shinkurt dimma	Spice, food
2.	Allium sativum	Nech shinkurt	Spice, food
3.	Artemisia abyssinica	Harit	spice
4.	Brassica carinata	Gomenzer	food
5.	Brassica nigra	Senafich	food
6.	Capsicum annum	Berbere	food
7.	Carica papaya	Papaya	food
8.	Citrus limon	Lomi	Food,
9.	Coffea Arabica	Buna	stimulant
10.	Cordia africana	Wanza	Timber, live fence, fire wood
11.	Coriandrum sativum	Denbelal	Food
12.	Croton macrostchys	Bakanisaa[bisana]	Fire wood,
13.	Cucurbita pepo	Duba	Food
14.	Echinops kebericho	Kerebicho	Smell
15.	Eucalyptus globulus	Bahirzaf Addii.	Construction, fire wood, live fence
16.	Guizotia abyssinica	Nug	Food
17.	Linum usitatissimum	Telba	Food

18.	<i>Lepidium sativum</i>	Feto	Food
19.	<i>Nigella sativa</i>	Tikur azmud	Food
20.	<i>Ocimum basilicum</i>	Zikakibe	Spice
21.	<i>Olea europaea</i> L. subsp. <i>cuspidata</i>	Weyra	Fire wood, consruction, farming tool
22.	<i>Otostegia integrifolia</i>	Tunjit	Fumigate
23.	<i>Prunus persica</i>	Kok	Food
24.	<i>Rhamnus prinoides</i>	Gesho	Bevarage
25.	<i>Ruta chalepensis</i>	Tenadam	Spice
26.	<i>Schinus molle</i>	Kundo berbere	Spice
27.	<i>Sesamum indicum</i>	Seli	Food
28.	<i>Trigonella foenum-Graecum</i>	[Sunqoo]Abish	Food
29.	<i>Thymus schimperi</i>	Tosign	Stimulant
30.	<i>Zingiber officinale</i>	Jinjiblaa	Spice